

$\beta(\tau_1, \tau_2)$ -CONTINUOUS MULTIFUNCTIONS ON BITOPOLOGICAL
SPACES

KAEWTA LAPROM


พหุบัณฑิต ชีวะ

A Thesis Submitted in Partial Fulfillment of Requirements for
Master of Science (Mathematics Education)

March 2020

Copyright of Mahasarakham University


ฟังก์ชันหลายค่าต่อเนื่อง $\beta(\tau_1, \tau_2)$ บนปริภูมิเชิงไมโทพอโลยี



เสนอต่อมหาวิทยาลัยมหาสารคาม เพื่อเป็นส่วนหนึ่งของการศึกษาตามหลักสูตร
ปริญญาวิทยาศาสตรมหาบัณฑิต สาขาวิชาคณิตศาสตร์ศึกษา

มีนาคม 2563

ลิขสิทธิ์เป็นของมหาวิทยาลัยมหาสารคาม



The examining committee has unanimously approved this thesis, submitted by Miss Kaewta Laprom, as a partial fulfillment of the requirements for the Master of Science in Mathematics Education at Mahasarakham University.

Examining Committee

.....	Chairman
(Asst. Prof. Supunnee Sompong, Ph.D.)	(Faculty graduate committee)
.....	Committee
(Asst. Prof. Chawalit Boonpok, Ph.D.)	(Advisor)
.....	Committee
(Asst. Prof. Chokchai Viriyapong, Ph.D.)	(Co-advisor)
.....	Committee
(Asst. Prof. Jeeranunt Khampakdee, Ph.D.)	(Faculty graduate committee)
.....	Committee
(Chalongchai Klanarong, Ph.D.)	(Faculty graduate committee)

Mahasarakham University has granted approval to accept this thesis as a partial fulfillment of the requirements for the Master of Science in Mathematics Education.

.....
(Prof. Pairot Pramual , Ph.D.)
Dean of the Faculty of Science

.....
(Assoc. Prof. Krit Chaimoon, Ph.D.)
Dean of Graduate School

ACKNOWLEDGEMENTS

Firstly, I would like to express my sincere gratitude to my advisor, Asst. Prof. Chawalit Boonpok, and co-advisor, Asst. Prof. Chokchai Viriyapong, for the continuous support of my thesis, and for their patience, motivation, and immense knowledge. Their guidance helped me in all the time of investigating my interesting theorems and writing of this thesis. I could not have imagined having a better advisor or mentor for my study.


Besides my advisor, I would like to thank the rest of my thesis committee: Asst. Prof. Supunee Sompong, Asst. Prof. Jeeranunt Khampakdee and Dr. Chalongchai Khanarong, not only for their insightful comments and encouragement but also for interesting questions.

My sincere thanks also go to my seniors, who answered my questions that were important to my study. Without their precious supports, it would not be possible to conduct this thesis.

I would like to thank my family including my parents and my brother for supporting me spiritually throughout writing this thesis and my life in general.

Lastly, I cannot have opportunity to be a graduate student without the support of the government unit called “The Promotion of Science and Mathematics Talented Teacher (PSMT)”, which provided me a scholarship until I graduated.

Kaewta Laprom




พหุ ประถมศึกษา

TITLE $\beta(\tau_1, \tau_2)$ -Continuous Multifunctions on Bitopological Spaces
CANDIDATE Kaewta Laprom
DEGREE Master of Science **MAJOR** Mathematics Education
ADVISORS Asst. Prof. Chawalit Boonpok, Ph.D.
 Asst. Prof. Chokchai Viriyapong, Ph.D.
UNIVERSITY Mahasarakham University **YEAR** 2020

ABSTRACT

The purpose of this paper is to introduce the concepts of $\beta(\tau_1, \tau_2)$ -continuous multifunctions, almost $\beta(\tau_1, \tau_2)$ -continuous multifunctions, and weakly $\beta(\tau_1, \tau_2)$ -continuous multifunctions. Moreover, some characterizations of these multifunctions are investigated. Besides, the relationships between upper and lower $\beta(\tau_1, \tau_2)$ -continuous multifunctions and the other types of continuity are also discussed.

Keywords : $\tau_1\tau_2$ - β -open, upper $\beta(\tau_1, \tau_2)$ -continuous multifunction, lower $\beta(\tau_1, \tau_2)$ -continuous multifunction




ชื่อเรื่อง	ฟังก์ชันหลายค่าต่อเนื่อง $\beta(\tau_1, \tau_2)$ บนปริภูมิเชิงไมโทพอโลยี
ผู้วิจัย	นางสาวแก้วตา หล้าพรหม
ปริญญา	วิทยาศาสตรมหาบัณฑิต สาขา คณิตศาสตร์ศึกษา
อาจารย์ที่ปรึกษา	ผู้ช่วยศาสตราจารย์ ดร. ชวลิต บุญปก ผู้ช่วยศาสตราจารย์ ดร.โชคชัย วิริยะพงษ์
มหาวิทยาลัย	มหาวิทยาลัยมหาสารคาม ปีที่พิมพ์ 2563

บทคัดย่อ

วัตถุประสงค์ของการวิจัยครั้งนี้เพื่อนำเสนอแนวคิดของฟังก์ชันหลายค่าต่อเนื่อง $\beta(\tau_1, \tau_2)$ ฟังก์ชันหลายค่าเกือบต่อเนื่อง $\beta(\tau_1, \tau_2)$ และฟังก์ชันหลายค่าต่อเนื่องอย่างอ่อน $\beta(\tau_1, \tau_2)$ อีกทั้งแสดงให้เห็นลักษณะของฟังก์ชันหลายค่าเหล่านี้ นอกจากนี้ยังกล่าวถึงความสัมพันธ์ระหว่างฟังก์ชันหลายค่าต่อเนื่อง $\beta(\tau_1, \tau_2)$ กับความต่อเนื่องประเภทอื่นๆ

คำสำคัญ : เซตเปิด $\tau_1\tau_2-\beta$, ฟังก์ชันหลายค่าต่อเนื่องบน $\beta(\tau_1, \tau_2)$, ฟังก์ชันหลายค่าต่อเนื่องล่าง $\beta(\tau_1, \tau_2)$



CONTENTS

	Page
Acknowledgements	i
Abstract in English	ii
Abstract in Thai	iii
Contents	iv
Chapter 1 Introduction	1
1.1 Background	1
1.2 Objective of the research	1
1.3 Research methodology	2
1.4 Scope of the study	2
Chapter 2 Preliminaries	3
2.1 Bitopological spaces	3
2.2 Multifunctions	12
Chapter 3 $\beta(\tau_1, \tau_2)$-continuous multifunctions	14
3.1 Characterizations of upper and lower $\beta(\tau_1, \tau_2)$ -continuous multifunctions	14
3.2 Characterizations of upper and lower almost $\beta(\tau_1, \tau_2)$ -continuous multifunctions	21
Chapter 4 Weakly $\beta(\tau_1, \tau_2)$-continuous multifunctions	27
4.1 Weakly $\beta(\tau_1, \tau_2)$ -continuous multifunctions	27
Chapter 5 Conclusions	35
5.1 Conclusions	35
5.2 Recommendations	39
References	40
Biography	42

CHAPTER 1

INTRODUCTION

1.1 Background

General topology is an important mathematical branch which is applied for many fields of applied sciences. Continuity is a basic concept for the study in topological spaces. Generalization of this concept by using weaker forms of open sets such as semi-open sets [1], preopen sets [2] and β -open sets [3] is one of the main research topics of general topology. In 1983, Monsef et al. [4] introduced the classes of β -open sets called semi-preopen sets by Andrijević in [3]; moreover, Monsef et al. [4] introduced almost β -continuous functions in topological spaces. From 1992 to 1993, the authors[5] obtained several characterizations of β -continuity and showed that almost quasi-continuity [6] investigated by Borsik and Dobos was equivalent to β -continuity. Therefore, in 1997, Nasef and Noiri [7] investigated fundamental characterizations of almost β -continuous functions. A year later, Popa and Noiri [8] investigated further characterizations of almost β -continuous functions. In 1992, Khedr et al.[9] generalized the notions of β -open sets and investigated β -continuous functions in bitopological spaces. Furthermore, in [10], [11], and [12] from 1996 to 2000, the authors extended these functions to multifunctions by introducing and characterizing the notions of β -continuous multifunctions, almost β -continuous multifunctions, and weakly β -continuous multifunctions in topological spaces.

Therefore, we are interested in defining upper and lower $\beta(\tau_1, \tau_2)$ -continuous multifunctions and investigating some characterizations of these multifunctions. Furthermore, almost $\beta(\tau_1, \tau_2)$ -continuous multifunctions and weakly $\beta(\tau_1, \tau_2)$ -continuous multifunctions were investigated.

1.2 Objective of the research

The purposes of the research are:

- 1.2.1 To define and investigate the characterizations of upper and lower $\beta(\tau_1, \tau_2)$ -continuous multifunctions.

1.2.2 To define and investigate the characterizations of upper and lower almost $\beta(\tau_1, \tau_2)$ -continuous multifunctions.

1.2.3 To define and investigate the characterizations of upper and lower weakly $\beta(\tau_1, \tau_2)$ -continuous multifunctions.

1.3 Research methodology

The research procedure of this thesis consists of the following steps:

1.3.1 Criticism and possible extension of the literature review.

1.3.2 To define and investigate the characterizations of upper and lower $\beta(\tau_1, \tau_2)$ -continuous multifunctions.


1.3.3 To define and investigate the characterizations of upper and lower almost $\beta(\tau_1, \tau_2)$ -continuous multifunctions.

1.3.4 To define and investigate the characterizations of upper and lower weakly $\beta(\tau_1, \tau_2)$ -continuous multifunctions.

1.3.5 To make the conclusions and do a complete report to offer Mahasarakham University.

1.4 Scope of the study

The scopes of the study are: defining and investigating the characterizations of upper and lower $\beta(\tau_1, \tau_2)$ -continuous multifunctions, the characterizations of upper and lower almost $\beta(\tau_1, \tau_2)$ -continuous multifunctions, and the characterizations of upper and lower weakly $\beta(\tau_1, \tau_2)$ -continuous multifunctions.



CHAPTER 2

PRELIMINARIES

In this chapter, we will give some definitions, notations, dealing with some preliminaries and some useful results that will be duplicated in later chapter.

2.1 Bitopological spaces

Definition 2.1.1. [13] Let $X \neq \emptyset$ and τ be a collection of subsets of X . Then, τ is called a topology on X if and only if τ satisfies the properties:

1. $\emptyset, X \in \tau$.
2. If $G_1, G_2 \in \tau$ then $G_1 \cap G_2 \in \tau$.
3. If $G_i \in \tau$ for all $i \in J$ then $\cup_{i \in J} G_i \in \tau$.

The pair (X, τ) is called a topological space. Sometimes this research, spaces (X, τ) (or simply X) always mean topological spaces.

Definition 2.1.2. [13] Let X be a topological space, and $A \subseteq X$. The *interior* of A is the set given by $\text{Int}(A) = \cup\{U \subseteq X : U \subseteq A \text{ and } U \text{ is open}\}$.

Theorem 2.1.3. [13] If X is a space and $A \subseteq X$, then the following are true:

1. $\text{Int}(A)$ is an open set.
2. $\text{Int}(A) \subseteq A$.
3. If $A \subseteq B \subseteq X$, then $\text{Int}(A) \subseteq \text{Int}(B)$.
4. If U is an open set with $U \subseteq A$, then $U \subseteq \text{Int}(A)$, that is $\text{Int}(A)$ is the largest open set contained in A .
5. For every $A \subseteq X$ and $B \subseteq X$, $\text{Int}(A \cap B) = \text{Int}(A) \cap \text{Int}(B)$.
6. For every $A \subseteq X$, A is open if and only if $A = \text{Int}(A)$.

Definition 2.1.4. [13] If A is a subset A of a topological space X , then the *closure* of A is the set given by $\text{Cl}(A) = \cap \{F \subseteq X : F \text{ is closed and } A \subseteq F\}$.

Theorem 2.1.5. [13] If X is a space and $A \subseteq X$, then the following are true:

1. $\text{Cl}(A)$ is an closed set.
2. $A \subseteq \text{Cl}(A)$.
3. If $A \subseteq B \subseteq X$, then $\text{Cl}(A) \subseteq \text{Cl}(B)$.
4. If F is an closed set with $A \subseteq F$, then $\text{Cl}(A) \subseteq F$, that is $\text{Cl}(A)$ is the smallest closed set containing in A .
5. For every $A \subseteq X$ and $B \subseteq X$, $\text{Cl}(A \cup B) = \text{Cl}(A) \cup \text{Cl}(B)$.
6. For every $A \subseteq X$, A is closed if and only if $A = \text{Cl}(A)$.

Proposition 2.1.6. [3] For a subset A of a topological space (X, τ) , the following properties hold:

- (1) $\text{Cl}(A) \cap G \subseteq \text{Cl}(A \cap G)$ for every open set G .
- (2) $\text{Int}(A \cup F) \subseteq \text{Int}(A) \cup F$ for every closed set F .

Definition 2.1.7. [15] A collection \mathcal{A} of subsets of a space X is said to *cover* X , or to be a *covering* of X , if the union of the element of \mathcal{A} is equal to X . It is called an *open covering* of X if its elements are open subsets of X .

In this investigation, we study on bitopological spaces. According to Kelly [14], a bitopological space (X, τ_1, τ_2) is a set X with two topologies, τ_1 and τ_2 on the space. Let A be a subsets of bitopological space (X, τ_1, τ_2) . The closure of A and the interior of A with τ_i are denoted by $\tau_i\text{-Cl}(A)$ and $\tau_i\text{-Int}(A)$, respectively, for $i = 1, 2$.

Definition 2.1.8. [16] Let (X, τ_1, τ_2) be a bitopological space and let $A \subseteq X$. Then, A is called (i, j) -*semi-open* if $A \subseteq j\text{Cl}(i\text{Int}(A))$, where $i \neq j = 1, 2$. The complement of (i, j) -semi-open set is (i, j) -*semi-closed*.

Definition 2.1.9. [17] Let (X, τ_1, τ_2) be a bitopological space and let $A \subseteq X$. Then, A is called (i, j) -regular-open if $A = i\text{Int}(j\text{Cl}(A))$, where $i \neq j = 1, 2$. The complement of (i, j) -regular-open set is (i, j) -regular-closed.

Definition 2.1.10. [18] Let (X, τ_1, τ_2) be a bitopological space and let $A \subseteq X$. Then, A is called (i, j) -preopen if $A \subseteq i\text{Int}(j\text{Cl}(A))$. The complement of (i, j) -preopen set is (i, j) -preclosed.

Definition 2.1.11. [9] Let (X, τ_1, τ_2) be a bitopological space and let $A \subseteq X$. Then, A is called (i, j) - β -open if $A \subseteq j\text{Cl}(i\text{Int}(j\text{Cl}(A)))$. The complement of (i, j) - β -open set is (i, j) - β -closed.

Note: In our research, we let $\tau_1\tau_2$ -semi-open, $\tau_1\tau_2$ -preopen, $\tau_1\tau_2$ - β -open and $\tau_1\tau_2$ -regular-open represent $(2, 1)$ -semi-open, $(1, 2)$ -preopen, $(2, 1)$ - β -open and $(1, 2)$ -regular-open respectively; moreover, we do complement of those in the similar way.

Example 2.1.12. Let $X = \{a, b, c\}$ with topologies $\tau_1 = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$ and $\tau_2 = \{\emptyset, \{a\}, \{a, b\}, X\}$.

1. It is easy to verify that there are five $\tau_1\tau_2$ -semi-open such as $\emptyset, \{a\}, \{a, b\}, \{a, c\}$, and X ; furthermore, five $\tau_1\tau_2$ -semi-closed sets on X are $\emptyset, \{b, c\}, \{c\}, \{b\}$, and X .
2. It is easy to verify that there are three $\tau_1\tau_2$ -regular-open such as $\emptyset, \{b\}$ and X ; furthermore, three $\tau_1\tau_2$ -regular-closed sets on X are $\emptyset, \{a, c\}$, and X .
3. It is easy to verify that the $\tau_1\tau_2$ -preopen on X are $\emptyset, \{a\}, \{b\}, \{a, b\}, \{a, c\}$ and X ; furthermore, six $\tau_1\tau_2$ -preclosed sets on X are $\emptyset, \{b, c\}, \{a, c\}, \{c\}, \{b\}$, and X .
4. It is easy to check that $\{a\}$ is a $\tau_1\tau_2$ - β -open set but $\{c\}$ is not a $\tau_1\tau_2$ - β -open set. Therefore, we know the complement of $\{a\}$ is $\tau_1\tau_2$ - β -closed set.

Definition 2.1.13. [19] Let (X, τ_1, τ_2) be a bitopological space and A be a subset of X . Then, $\tau_1\tau_2$ -semi-closure of A denoted by $\tau_1\tau_2\text{-sCl}(A)$ is defined as

$$\tau_1\tau_2\text{-sCl}(A) = \cap\{F \subseteq X : F \text{ is } \tau_1\tau_2\text{-semi-closed in } X \text{ and } A \subseteq F\}.$$

Definition 2.1.14. [9] Let (X, τ_1, τ_2) be a bitopological space and A be a subset of X . Then, $\tau_1\tau_2$ -preclosure of A denoted by $\tau_1\tau_2\text{-pCl}(A)$ is defined as

$$\tau_1\tau_2\text{-pCl}(A) = \cap\{F \subseteq X : F \text{ is } \tau_1\tau_2\text{-preclosed in } X \text{ and } A \subseteq F\}.$$

Definition 2.1.15. [9] Let (X, τ_1, τ_2) be a bitopological space and A be a subset of X . Then, $\tau_1\tau_2$ - β -closure of A denoted by $\tau_1\tau_2\text{-}\beta\text{Cl}(A)$ is defined as

$$\tau_1\tau_2\text{-}\beta\text{Cl}(A) = \cap\{F \subseteq X : F \text{ is } \tau_1\tau_2\text{-}\beta\text{-closed in } X \text{ and } A \subseteq F\}.$$

Example 2.1.16. Let $X = \{a, b, c\}$ with topologies $\tau_1 = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$ and $\tau_2 = \{\emptyset, \{a\}, \{a, b\}, X\}$. Let $A = \{c\}$. Then, (X, τ_1, τ_2) is a bitopological space. In example 2.1.12, $\tau_1\tau_2$ -semi-closed in X are $\emptyset, \{b, c\}, \{c\}, \{b\}$, and X . Clearly, $A \subseteq \{b, c\}$ and $A \subseteq \{c\}$, then $\tau_1\tau_2\text{-sCl}(A) = \{b, c\} \cap \{c\} = \{c\}$. Moreover, in example 2.1.12, $\tau_1\tau_2$ -preclosed in X are $\emptyset, \{b, c\}, \{a, c\}, \{b\}, \{c\}$, and X . Clearly, $A \subseteq \{b, c\}$, $A \subseteq \{c\}$ and $A \subseteq \{a, c\}$. Hence, $\tau_1\tau_2\text{-pCl}(A) = \{b, c\} \cap \{c\} \cap \{a, c\} = \{c\}$. Moreover, it is easy to check that $\tau_1\tau_2\text{-}\beta\text{Cl}(A) = \{c\}$.

Definition 2.1.17. [19] Let (X, τ_1, τ_2) be a bitopological space and A be a subset of X . Then, $\tau_1\tau_2$ -semi-interior of A denoted by $\tau_1\tau_2\text{-sInt}(A)$ is defined as

$$\tau_1\tau_2\text{-sInt}(A) = \cup\{G \subseteq X : G \text{ is } \tau_1\tau_2\text{-semi-open in } X \text{ and } G \subseteq A\}.$$

Definition 2.1.18. [20] Let (X, τ_1, τ_2) be a bitopological space and A be a subset of X . Then, $\tau_1\tau_2$ -preinterior of A denoted by $\tau_1\tau_2\text{-pInt}(A)$ is defined as

$$\tau_1\tau_2\text{-pInt}(A) = \cup\{G \subseteq X : G \text{ is } \tau_1\tau_2\text{-preopen in } X \text{ and } G \subseteq A\}.$$

Definition 2.1.19. [9] Let (X, τ_1, τ_2) be a bitopological space and A be a subset of X . Then, $\tau_1\tau_2$ - β -interior of A denoted by $\tau_1\tau_2\text{-}\beta\text{Int}(A)$ is defined as

$$\tau_1\tau_2\text{-}\beta\text{Int}(A) = \cup\{G \subseteq X : G \text{ is } \tau_1\tau_2\text{-}\beta\text{-open in } X \text{ and } G \subseteq A\}.$$

Example 2.1.20. Let $X = \{a, b, c\}$ with topologies $\tau_1 = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$ and $\tau_2 = \{\emptyset, \{a\}, \{a, b\}, X\}$. Let $A = \{a, c\}$. Then, (X, τ_1, τ_2) is a bitopological space.

In example 2.1.12, $\tau_1\tau_2$ -semi-open in X are $\emptyset, \{a\}, \{a, b\}, \{a, c\}$, and X . Clearly, $\emptyset \subseteq A$, $\{a\} \subseteq A$ and $\{a, c\} \subseteq A$, then $\tau_1\tau_2\text{-sInt}(A) = \emptyset \cup \{a\} \cup \{a, c\} = \{a, c\}$.

Moreover, in example 2.1.12 is that $\tau_1\tau_2$ -preopen in X are $\emptyset, \{a\}, \{b\}, \{a, c\}, \{a, b\}$, and X . Clearly, $\emptyset \subseteq A$, $\{a\} \subseteq A$ and $\{a, c\} \subseteq A$. Hence, $\tau_1\tau_2\text{-pInt}(A) = \emptyset \cup \{a\} \cup \{a, c\} = \{a, c\}$. Moreover, it is easy to check that $\tau_1\tau_2\text{-}\beta\text{Int}(A) = \{a, c\}$.

Definition 2.1.21. [21] Let (X, τ_1, τ_2) be a bitopological space and let $A \subseteq X$. Then, A is called $\tau_1\tau_2$ -open if $A = \tau_1\text{-Int}(\tau_2\text{-Int}(A))$. The complement of $\tau_1\tau_2$ -open set is $\tau_1\tau_2$ -closed.

Example 2.1.22. Let $X = \{a, b, c\}$ with topologies $\tau_1 = \{\emptyset, \{b\}, \{c\}, \{b, c\}, X\}$ and $\tau_2 = \{\emptyset, \{a\}, \{b, c\}, X\}$. It is easy to check that $\{b, c\}$ is a $\tau_1\tau_2$ -open set but $\{c\}$ is not a $\tau_1\tau_2$ -open set. Therefore, we know the complement of $\{b, c\}$ is $\tau_1\tau_2$ -closed set.

Definition 2.1.23. [21] Let (X, τ_1, τ_2) be a bitopological space and A be a subset of X . Then, $\tau_1\tau_2$ -closure of A denoted by $\tau_1\tau_2\text{-Cl}(A)$ is defined as

$$\tau_1\tau_2\text{-Cl}(A) = \cap\{F \subseteq X : F \text{ is } \tau_1\tau_2\text{-closed in } X \text{ and } A \subseteq F\}.$$

Example 2.1.24. Let $X = \{a, b, c\}$ with topologies $\tau_1 = \{\emptyset, \{b\}, \{c\}, \{b, c\}, X\}$ and $\tau_2 = \{\emptyset, \{a\}, \{b, c\}, X\}$. Let $A = \{a\}$. Then, (X, τ_1, τ_2) is a bitopological space. It is easy to verify that $\tau_1\tau_2$ -closed in X are $\emptyset, \{a\}$, and X . Clearly, $A \subseteq \{a\}$ and $A \subseteq X$, then $\tau_1\tau_2\text{-Cl}(A) = \{a\} \cap X = \{a\}$.

Definition 2.1.25. [21] Let (X, τ_1, τ_2) be a bitopological space and A be a subset of X . Then, $\tau_1\tau_2$ -interior of A denoted by $\tau_1\tau_2\text{-Int}(A)$ is defined as

$$\tau_1\tau_2\text{-Int}(A) = \cup\{G \subseteq X : G \text{ is } \tau_1\tau_2\text{-open in } X \text{ and } G \subseteq A\}.$$

Example 2.1.26. Let $X = \{a, b, c\}$ with topologies $\tau_1 = \{\emptyset, \{b\}, \{c\}, \{b, c\}, X\}$ and $\tau_2 = \{\emptyset, \{a\}, \{b, c\}, X\}$. Let $A = \{b, c\}$. Then, (X, τ_1, τ_2) is a bitopological space. It is easy to verify that $\tau_1\tau_2$ -open in X are $\emptyset, \{b, c\}$, and X . Clearly, $\emptyset \subseteq A$ and $\{b, c\} \subseteq A$, then $\tau_1\tau_2\text{-Int}(A) = \emptyset \cup \{b, c\} = \{b, c\}$.

Definition 2.1.27. [21] A set N of a bitopological space (X, τ_1, τ_2) is said to be $\tau_1\tau_2$ -neighbourhood of $x \in X$ if there exists a $\tau_1\tau_2$ -open set V of (X, τ_1, τ_2) such that $x \in V \subseteq N$.

Example 2.1.28. Let $X = \{a, b, c\}$ with topologies $\tau_1 = \{\emptyset, \{b\}, \{c\}, \{b, c\}, X\}$ and $\tau_2 = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{b, c\}, X\}$. Let $N = \{a, b\}$.

Then, (X, τ_1, τ_2) is a bitopological space. It is easy to verify that $\tau_1\tau_2$ -open in X are $\emptyset, \{b\}, \{b, c\}$, and X . Clearly, $b \in \{b\} \subseteq N$, then N is $\tau_1\tau_2$ -neighbourhood of $b \in X$.

Definition 2.1.29. A set N of a bitopological space (X, τ_1, τ_2) is said to be $\tau_1\tau_2$ -preneighbourhood of $x \in X$ if there exists a $\tau_1\tau_2$ -preopen set V of (X, τ_1, τ_2) such that $x \in V \subseteq N$.

Example 2.1.30. Let $X = \{a, b, c\}$ with topologies $\tau_1 = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$, $\tau_2 = \{\emptyset, \{a\}, \{a, b\}, X\}$ and $N = \{a, c\}$, then (X, τ_1, τ_2) is the bitopological space. In example 2.1.12, the $\tau_1\tau_2$ -preopen on X are $\emptyset, \{a\}, \{b\}, \{a, b\}, \{a, c\}$ and X . Clearly, $a \in \{a\} \subseteq N$, then N is $\tau_1\tau_2$ -preneighbourhood of $a \in X$.

Proposition 2.1.31. [21] Let A and B be subsets of a bitopological space (X, τ_1, τ_2) . For the $\tau_1\tau_2$ -closure, the following properties hold:

- (1) $A \subseteq \tau_1\tau_2\text{-Cl}(A)$ and $\tau_1\tau_2\text{-Cl}(\tau_1\tau_2\text{-Cl}(A)) = \tau_1\tau_2\text{-Cl}(A)$.
- (2) If $A \subseteq B$, then $\tau_1\tau_2\text{-Cl}(A) \subseteq \tau_1\tau_2\text{-Cl}(B)$.
- (3) $\tau_1\tau_2\text{-Cl}(A)$ is $\tau_1\tau_2$ -closed.
- (4) A is $\tau_1\tau_2$ -closed if and only if $A = \tau_1\tau_2\text{-Cl}(A)$.
- (5) $\tau_1\tau_2\text{-Cl}(X - A) = X - \tau_1\tau_2\text{-Int}(A)$.

Proposition 2.1.32. Let (X, τ_1, τ_2) be a bitopological space. If A is $\tau_1\tau_2$ -semi-open and B is $\tau_1\tau_2$ -open in X , then $A \cap B$ is $\tau_1\tau_2$ -semi-open.

Proof. Suppose that A is $\tau_1\tau_2$ -semi-open and B is $\tau_1\tau_2$ -open in X .

Then, $A \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(A))$ and $B = \tau_1\text{-Int}(B) = \tau_2\text{-Int}(B)$. Therefore, we obtain $A \cap B \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(A)) \cap B$. By Proposition 2.1.6(1), we have

$$A \cap B \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(A)) \cap B \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(A) \cap B) = \tau_1\text{-Cl}(\tau_2\text{-Int}(A \cap B)).$$

Hence, $A \cap B$ is $\tau_1\tau_2$ -semi-open. □

Theorem 2.1.33. [9] Let (X, τ_1, τ_2) be a bitopological. Let $\{A_\gamma | \gamma \in \nabla\}$ be a family of subsets of X . The following properties are hold:

- (1) If A_γ is $\tau_1\tau_2$ -semi-open for each $\gamma \in \nabla$, then $\cup_{\gamma \in \nabla} A_\gamma$ is $\tau_1\tau_2$ -semi-open.
- (2) If A_γ is $\tau_1\tau_2$ -semi-closed for each $\gamma \in \nabla$, then $\cap_{\gamma \in \nabla} A_\gamma$ is $\tau_1\tau_2$ -semi-closed.

Proof. 1. Suppose that A_γ is $\tau_1\tau_2$ -semi-open for each $\gamma \in \nabla$. Then, we have

$$A_\gamma \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(A_\gamma)) \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(\cup_{\gamma \in \nabla} A_\gamma)), \text{ and hence}$$

$$\cup_{\gamma \in \nabla} A_\gamma \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(\cup_{\gamma \in \nabla} A_\gamma)). \text{ This shows that } \cup_{\gamma \in \nabla} A_\gamma \text{ is } \tau_1\tau_2\text{-semi-open.}$$

2. Suppose that A_γ is $\tau_1\tau_2$ -semi-closed for each $\gamma \in \nabla$. Then, we have

$$X - A_\gamma \text{ is } \tau_1\tau_2\text{-semi-open and } X - \cap_{\gamma \in \nabla} A_\gamma = \cup_{\gamma \in \nabla} (X - A_\gamma). \text{ Therefore, by}$$

$$(1), X - \cap_{\gamma \in \nabla} A_\gamma \text{ is } \tau_1\tau_2\text{-semi-open, and hence } \cap_{\gamma \in \nabla} A_\gamma \text{ is } \tau_1\tau_2\text{-semi-closed.}$$

□

Proposition 2.1.34. For a subset A of a bitopological space (X, τ_1, τ_2) , the following properties hold:

- (1) $\tau_1\tau_2\text{-sInt}(A)$ is $\tau_1\tau_2$ -semi-open.
- (2) $\tau_1\tau_2\text{-sCl}(A)$ is $\tau_1\tau_2$ -semi-closed.
- (3) A is $\tau_1\tau_2$ -semi-open if and only if $A = \tau_1\tau_2\text{-sInt}(A)$.
- (4) A is $\tau_1\tau_2$ -semi-closed if and only if $A = \tau_1\tau_2\text{-sCl}(A)$.

Proof. (1) and (2) follows from Proposition 2.1.33 (3) and (4) follows from (1) and (2). □

Proposition 2.1.35. For a subset A of a bitopological space (X, τ_1, τ_2) , $x \in \tau_1\tau_2\text{-sCl}(A)$ if and only if $U \cap A \neq \emptyset$ for every $\tau_1\tau_2$ -semi-open set U containing x .

Proof. Let $x \in \tau_1\tau_2\text{-sCl}(A)$. We shall show that $U \cap A \neq \emptyset$ for every $\tau_1\tau_2$ -semi-open set U containing x . Suppose that $U \cap A = \emptyset$ for some $\tau_1\tau_2$ -semi-open set U containing x . Then, $A \subseteq X - U$ and $X - U$ is $\tau_1\tau_2$ -semi-closed. Since $x \in \tau_1\tau_2\text{-sCl}(A)$, we have $x \in \tau_1\tau_2\text{-sCl}(X - U) = X - U$; hence $x \notin U$, which is a contradiction that $x \in U$. Therefore, $U \cap A \neq \emptyset$.

Conversely, we assume that $U \cap A \neq \emptyset$ for every $\tau_1\tau_2$ -semi-open set U containing x . We shall show that $x \in \tau_1\tau_2\text{-sCl}(A)$. Suppose that $x \notin \tau_1\tau_2\text{-sCl}(A)$. Then, there exists a $\tau_1\tau_2$ -semi-closed set F such that $A \subseteq F$ and $x \notin F$. Therefore, we obtain

$X - F$ is a $\tau_1\tau_2$ -semi-open set containing x such that $(X - F) \cap A = \emptyset$. This a contradiction to $U \cap A \neq \emptyset$; hence $x \in \tau_1\tau_2\text{-sCl}(A)$. \square

Proposition 2.1.36. For a subset A of a bitopological space (X, τ_1, τ_2) , the following properties hold:

$$(1) X - \tau_1\tau_2\text{-sCl}(A) = \tau_1\tau_2\text{-sInt}(X - A).$$

$$(2) X - \tau_1\tau_2\text{-sInt}(A) = \tau_1\tau_2\text{-sCl}(X - A).$$

Proof. (1) Let $x \in X - \tau_1\tau_2\text{-sCl}(A)$. Then, $x \notin \tau_1\tau_2\text{-sCl}(A)$. By Proposition 2.1.35, there exists a $\tau_1\tau_2$ -semi-open set V containing x such that $V \cap A = \emptyset$. Then, $V \subseteq X - A$, and hence $x \in \tau_1\tau_2\text{-sInt}(X - A)$. This shows that $X - \tau_1\tau_2\text{-sCl}(A) \subseteq \tau_1\tau_2\text{-sInt}(X - A)$.

Let $x \in \tau_1\tau_2\text{-sInt}(X - A)$. Then, there exists a $\tau_1\tau_2$ -semi-open set V containing x such that $V \subseteq X - A$, and hence $V \cap A = \emptyset$. By Proposition 2.1.35, we have $x \notin \tau_1\tau_2\text{-sCl}(A)$; hence $x \in X - \tau_1\tau_2\text{-sCl}(A)$. Therefore,

$$\tau_1\tau_2\text{-sInt}(X - A) \subseteq X - \tau_1\tau_2\text{-sCl}(A).$$

Consequently, we obtain $X - \tau_1\tau_2\text{-sCl}(A) = \tau_1\tau_2\text{-sInt}(X - A)$.

(2) This follow from (1). \square

Proposition 2.1.37. For a subset A of a bitopological space (X, τ_1, τ_2) , the following properties hold:

$$(1) \tau_1\tau_2\text{-sCl}(A) = \tau_1\text{-Int}(\tau_2\text{-Cl}(A)) \cup A.$$

$$(2) \text{ If } A \text{ is } \tau_1\text{-open in } X, \text{ then } \tau_1\tau_2\text{-sCl}(A) = \tau_1\text{-Int}(\tau_2\text{-Cl}(A)).$$

Proof. (1) Since $\tau_1\tau_2\text{-sCl}(A)$ is $\tau_1\tau_2$ -semi-closed, we have

$$\tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\tau_2\text{-sCl}(A))) \subseteq \tau_1\tau_2\text{-sCl}(A).$$

Thus, $\tau_1\text{-Int}(\tau_2\text{-Cl}(A)) \subseteq \tau_1\tau_2\text{-sCl}(A)$. Hence, $\tau_1\text{-Int}(\tau_2\text{-Cl}(A)) \cup A \subseteq \tau_1\tau_2\text{-sCl}(A)$.

To establish the opposite inclusion, we observe that

$$\begin{aligned} \tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\text{-Int}(\tau_2\text{-Cl}(A)) \cup A)) &\subseteq \tau_1\text{-Int}(\tau_2\text{-Cl}(A) \cup A) \\ &= \tau_1\text{-Int}(\tau_2\text{-Cl}(A)). \end{aligned}$$

Therefore,

$$\tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\text{-Int}(\tau_2\text{-Cl}(A)) \cup A)) \subseteq \tau_1\text{-Int}(\tau_2\text{-Cl}(A)) \subseteq \tau_1\text{-Int}(\tau_2\text{-Cl}(A)) \cup A.$$

Hence, $\tau_1\text{-Int}(\tau_2\text{-Cl}(A)) \cup A$ is $\tau_1\tau_2$ -semi-closed. Then,

$$\tau_1\tau_2\text{-sCl}(A) \subseteq \tau_1\text{-Int}(\tau_2\text{-Cl}(A)) \cup A.$$

Consequently, we obtain $\tau_1\tau_2\text{-sCl}(A) = \tau_1\text{-Int}(\tau_2\text{-Cl}(A)) \cup A$.

(2) Let A be a τ_1 -open set, then $A = \tau_1\text{-Int}(A) \subseteq \tau_1\text{-Int}(\tau_2\text{-Cl}(A))$. Therefore, by (1), we have $\tau_1\tau_2\text{-sCl}(A) = \tau_1\text{-Int}(\tau_2\text{-Cl}(A))$. \square

Proposition 2.1.38. Let (X, τ_1, τ_2) be a bitopological space and $\{A_\gamma \mid \gamma \in \Gamma\}$ a family of subsets of X . The following properties hold:

- (1) If A_γ is $\tau_1\tau_2$ - β -open for each $\gamma \in \Gamma$, then $\bigcup_{\gamma \in \Gamma} A_\gamma$ is $\tau_1\tau_2$ - β -open.
- (2) If A_γ is $\tau_1\tau_2$ - β -closed for each $\gamma \in \Gamma$, then $\bigcap_{\gamma \in \Gamma} A_\gamma$ is $\tau_1\tau_2$ - β -closed.

Proof. 1. The proof follows from Theorem 3.2 of [9].

2. The proof follows from Lemma 2.1 of [22]. \square

Proposition 2.1.39. [22] For a subset A of a bitopological space (X, τ_1, τ_2) , the following properties hold:

- (1) $\tau_1\tau_2\text{-}\beta\text{Int}(A)$ is $\tau_1\tau_2$ - β -open.
- (2) $\tau_1\tau_2\text{-}\beta\text{Cl}(A)$ is $\tau_1\tau_2$ - β -closed.
- (3) A is $\tau_1\tau_2$ - β -open if and only if $A = \tau_1\tau_2\text{-}\beta\text{Int}(A)$.
- (4) A is $\tau_1\tau_2$ - β -closed if and only if $A = \tau_1\tau_2\text{-}\beta\text{Cl}(A)$.

Proposition 2.1.40. [22] For a subset A of a bitopological space (X, τ_1, τ_2) , $x \in \tau_1\tau_2\text{-}\beta\text{Cl}(A)$ if and only if $U \cap A \neq \emptyset$ for every $\tau_1\tau_2$ - β -open set U containing x .

Proposition 2.1.41. [22] For a subset A of a bitopological space (X, τ_1, τ_2) , the following properties hold:

$$(1) X - \tau_1\tau_2\text{-}\beta\text{Cl}(A) = \tau_1\tau_2\text{-}\beta\text{Int}(X - A).$$

$$(2) X - \tau_1\tau_2\text{-}\beta\text{Int}(A) = \tau_1\tau_2\text{-}\beta\text{Cl}(X - A).$$

2.2 Multifunctions

A multifunction $F : X \rightarrow Y$ is a point-to-set correspondence from X to Y .

We always assume that $F(x) \neq \emptyset$ for all $x \in X$. We shall denote the upper and lower inverse [23] of a set B of Y by $F^+(B)$ and $F^-(B)$, respectively, that is $F^+(B) = \{x \in X | F(x) \subseteq B\}$ and $F^-(B) = \{x \in X | F(x) \cap B \neq \emptyset\}$.

Example 2.2.1. Let $X = \{1, 2, 3\}$ and $Y = \{a, b, c, d, e\}$. A multifunction $F : X \rightarrow Y$ is defined as follows : $F(1) = \{c\}$, $F(2) = \{b, d\}$ and $F(3) = \{a, e\}$. Then, $F^+(\{a, b, c, d\}) = \{1, 2\}$ and $F^-(\{a, b, c, d\}) = \{1, 2, 3\}$. Moreover, it is easy to check that $F^-(\{a\}) = \{3\}$.

Definition 2.2.2. [23] A multifunction $F : X \rightarrow Y$ and $A \subseteq X$, then *direct image* of A under the multifunction F is the set $F(A) = \cup_{x \in A} F(x)$.

Example 2.2.3. Let $X = \{1, 2, 3\}$ and $Y = \{a, b, c, d, e\}$. A multifunction $F : X \rightarrow Y$ is defined as follows : $F(1) = \{c\}$, $F(2) = \{b, d\}$ and $F(3) = \{a, e\}$.

Therefore, if $A = \{1, 2\} \subseteq X$, then $F(A) = F(1) \cup F(2) = \{c\} \cup \{b, d\} = \{b, c, d\}$.

Lemma 2.2.4. Let $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ be a multifunction and $U \subseteq X, V \subseteq Y$. Then, the following properties hold:

- (1) If $F(U) \subseteq V$, then $U \subseteq F^+(V)$.
- (2) If $F(z) \cap V \neq \emptyset$ for every $z \in U$, then $U \subseteq F^-(V)$.
- (3) If $U \subseteq V$ then $F^+(U) \subseteq F^+(V)$.
- (4) If $U \subseteq V$ then $F^-(U) \subseteq F^-(V)$.
- (5) $X - F^-(V) = F^+(Y - V)$.


Proof. (1) Let $x \in U$. Since $\cup_{x \in U} F(x) = F(U) \subseteq V$, $F(x) \subseteq F(U) \subseteq V$. Thus, $F(x) \subseteq V$; hence $x \in F^+(V)$. This shows that $U \subseteq F^+(V)$.

(2) Suppose that $F(z) \cap V \neq \emptyset$ for every $z \in U$. Let $x \in U$. Therefore, $F(x) \cap V \neq \emptyset$. Hence, $x \in F^-(V)$. This shows that $U \subseteq F^-(V)$.

(3) Let $x \in F^+(U)$. Therefore, $F(x) \subseteq U$. Since $U \subseteq V$, $F(x) \subseteq V$. Thus, $x \in F^+(V)$. This shows that $F^+(U) \subseteq F^+(V)$.

(4) Let $x \in F^-(U)$. Therefore, $F(x) \cap U \neq \emptyset$. Since $U \subseteq V$, $F(x) \cap V \neq \emptyset$. Thus, $x \in F^-(V)$. This shows that $F^-(U) \subseteq F^-(V)$.

(5) Let $x \in X - F^-(V)$. Then, $x \notin F^-(V)$. Therefore, $F(x) \cap V = \emptyset$. Thus, $F(x) \subseteq Y - V$. Hence, $x \in F^+(Y - V)$. This implies that $X - F^-(V) \subseteq F^+(Y - V)$. On the other hand, suppose that $x \in F^+(Y - V)$, so $F(x) \subseteq Y - V$. Therefore, $F(x) \cap V = \emptyset$. Then, we obtain $x \notin F^-(V)$. Thus, $x \in X - F^-(V)$; hence, $F^+(Y - V) \subseteq X - F^-(V)$. Consequently, $X - F^-(V) = F^+(Y - V)$. \square



CHAPTER 3

$\beta(\tau_1, \tau_2)$ -CONTINUOUS MULTIFUNCTIONS

3.1 Characterizations of upper and lower $\beta(\tau_1, \tau_2)$ -continuous multifunctions

In this section, we introduce the notions of upper and lower $\beta(\tau_1, \tau_2)$ -continuous multifunctions, and investigate some characterizations of these multifunctions.

Definition 3.1.1. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is said to be:

- (1) *upper $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(U) \subseteq V$;
- (2) *lower $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y such that $F(x) \cap V \neq \emptyset$, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(z) \cap V \neq \emptyset$ for every $z \in U$;
- (3) *upper (resp. lower) $\beta(\tau_1, \tau_2)$ -continuous* if F has this property at each point of X .

Example 3.1.2. Let $X = \{a, b, c\}$ with topologies $\tau_1 = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$ and $\tau_2 = \{\emptyset, \{a\}, \{a, b\}, X\}$. Let $Y = \{1, 2, 3, 4, 5\}$ with topologies $\sigma_1 = \{\emptyset, \{1\}, \{2, 3, 4, 5\}, Y\}$ and $\sigma_2 = \{\emptyset, \{2\}, \{3\}, \{2, 3\}, \{2, 3, 4, 5\}, Y\}$.

A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is defined as follow: $F(a) = \{2, 3\}$, $F(b) = \{1, 2\}$, $F(c) = \{1, 4, 5\}$. Then, F is upper and lower $\beta(\tau_1, \tau_2)$ -continuous.

Theorem 3.1.3. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is upper $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(V))$ for every $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$.

Proof. Let V be a $\sigma_1\sigma_2$ -open set containing $F(x)$. Consequently, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(U) \subseteq V$. Therefore, $x \in U \subseteq F^+(V)$. Since U is $\tau_1\tau_2$ - β -open, we have $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(V))$.

Conversely, let V be a $\sigma_1\sigma_2$ -open set containing $F(x)$. By the hypothesis, $x \in \tau_1\tau_2$ - $\beta\text{Int}(F^+(V))$. There exists a $\tau_1\tau_2$ - β -open set G containing x such that $G \subseteq F^+(V)$; hence, $F(G) \subseteq V$. This shows that F is upper $\beta(\tau_1, \tau_2)$ -continuous at x . \square

Theorem 3.1.4. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2$ - $\beta\text{Int}(F^-(V))$ for every $\sigma_1\sigma_2$ -open set V of Y such that $F(x) \cap V \neq \emptyset$.

Proof. The proof is similar to that of Theorem 3.1.3. \square

Theorem 3.1.5. For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is upper $\beta(\tau_1, \tau_2)$ -continuous;
- (2) $F^+(V)$ is $\tau_1\tau_2$ - β -open in X for every $\sigma_1\sigma_2$ -open set V of Y ;
- (3) $F^-(K)$ is $\tau_1\tau_2$ - β -closed in X for every $\sigma_1\sigma_2$ -closed set K of Y ;
- (4) $\tau_1\tau_2$ - $\beta\text{Cl}(F^-(B)) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B of Y ;
- (5) $\tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\text{-Int}(F^-(B)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B of Y .

Proof. (2) \Rightarrow (1): Let $x \in X$ and V be a $\sigma_1\sigma_2$ -open set containing $F(x)$. By (2), $F^+(V)$ is a $\tau_1\tau_2$ - β -open set containing x . Putting $U = F^+(V)$, we obtain U is a $\tau_1\tau_2$ - β -open set containing x such that $F(U) \subseteq V$. This shows that F is upper $\beta(\tau_1, \tau_2)$ -continuous.

(1) \Rightarrow (2): Let V be a $\sigma_1\sigma_2$ -open set of Y and $x \in F^+(V)$. By 2.2.4, $F(x) \subseteq V$, then there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(U) \subseteq V$. Consequently, we obtain $x \in U \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(\tau_1\text{-Cl}(U))) \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(\tau_1\text{-Cl}(F^+(V))))$. Thus, $F^+(V) \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(\tau_1\text{-Cl}(F^+(V))))$. This shows $F^+(V)$ is $\tau_1\tau_2$ - β -open in X .

(2) \Rightarrow (3): This follows from the fact that $F^+(Y - B) = X - F^-(B)$ for every subset B of Y .

(3) \Rightarrow (4): For each subset B of Y , $\sigma_1\sigma_2\text{-Cl}(B)$ is $\sigma_1\sigma_2$ -closed in Y . By (3), $F^-(\sigma_1\sigma_2\text{-Cl}(B))$ is $\tau_1\tau_2$ - β -closed in X ; therefore, $\tau_1\tau_2$ - $\beta\text{Cl}(F^-(B)) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(B))$.

(4) \Rightarrow (5): Let B be a subset of Y . By Proposition 2.1.39(2), we obtain

$$\tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\text{-Int}(F^-(B)))) \subseteq \tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\text{-Int}(\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(B)))))$$

$$\subseteq \tau_1\tau_2\text{-}\beta\text{Cl}(F^-(B)).$$

Consequently, $\tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\text{-Int}(F^-(B)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(B))$ by (4).

(5) \Rightarrow (2): Let V be a $\sigma_1\sigma_2$ -open set of Y , so $Y - V$ is $\sigma_1\sigma_2$ -closed in Y . By (5) and Lemma 2.2.4,

$$\begin{aligned} X - \tau_1\text{-Cl}(\tau_2\text{-Int}(\tau_1\text{-Cl}(F^+(V)))) &= \tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\text{-Int}(X - F^+(V)))) \\ &= \tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\text{-Int}(F^-(Y - V)))) \\ &\subseteq F^-(\sigma_1\sigma_2\text{-Cl}(Y - V)) \\ &= X - F^+(\sigma_1\sigma_2\text{-Int}(V)) \\ &\subseteq X - F^+(V). \end{aligned}$$

Therefore, we obtain $F^+(V) \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(\tau_1\text{-Cl}(F^+(V))))$, and hence $F^+(V)$ is $\tau_1\tau_2\text{-}\beta$ -open in X . \square

Theorem 3.1.6. For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is lower $\beta(\tau_1, \tau_2)$ -continuous;
- (2) $F^-(V)$ is $\tau_1\tau_2\text{-}\beta$ -open in X for every $\sigma_1\sigma_2$ -open set V of Y ;
- (3) $F^+(K)$ is $\tau_1\tau_2\text{-}\beta$ -closed in X for every $\sigma_1\sigma_2$ -closed set K of Y ;
- (4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(B)) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B of Y ;
- (5) $\tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\text{-Int}(F^+(B)))) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B of Y .

Proof. It is shown similarly to the proof of Theorem 3.1.5 that the statements (1), (2), (3), (4) and (5) are equivalent. \square

Definition 3.1.7. [24] Let (X, τ_1, τ_2) be bitopological space. A covering \mathcal{B} is called a refinement of covering \mathcal{U} if every $\tau_1\tau_2$ -open set of \mathcal{B} is contained in some $\tau_1\tau_2$ -open set of \mathcal{U} .

Definition 3.1.8. [21] A collection \mathfrak{U} of subsets of a bitopological space (X, τ_1, τ_2) is said to be $\tau_1\tau_2$ -locally finite if every $x \in X$ has a $\tau_1\tau_2$ -neighborhood which intersects only finitely many elements of \mathfrak{U} .

Definition 3.1.9. [21] A subset A of a bitopological space (X, τ_1, τ_2) is said to be:

- (1) $\tau_1\tau_2$ -paracompact if every cover of A by $\tau_1\tau_2$ -open sets of X is refined by a cover of A which consists of $\tau_1\tau_2$ -open sets of X and is $\tau_1\tau_2$ -locally finite in X ;
- (2) $\tau_1\tau_2$ -regular if for each $x \in A$ and each $\tau_1\tau_2$ -open set U of X containing x , there exists a $\tau_1\tau_2$ -open set V of X such that $x \in V \subseteq \tau_1\tau_2\text{-Cl}(V) \subseteq U$.

Lemma 3.1.10. [21] If A is a $\tau_1\tau_2$ -regular $\tau_1\tau_2$ -paracompact set of a bitopological space (X, τ_1, τ_2) and U is a $\tau_1\tau_2$ -open neighbourhood of A , then there exists a $\tau_1\tau_2$ -open set V of X such that $A \subseteq V \subseteq \tau_1\tau_2\text{-Cl}(V) \subseteq U$.

Definition 3.1.11. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is called *punctually* (τ_1, τ_2) -paracompact (resp. *punctually* (τ_1, τ_2) -regular) if for each $x \in X$, $F(x)$ is $\tau_1\tau_2$ -paracompact (resp. $\tau_1\tau_2$ -regular).

For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, by

$$\beta\text{Cl}F_{\otimes} : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2),$$

we denote the multifunction defined as follows: $\beta\text{Cl}F_{\otimes}(x) = \sigma_1\sigma_2\text{-}\beta\text{Cl}(F(x))$ for each $x \in X$.

Example 3.1.12. Let $X = \{1, 2, 3\}$ with topologies

$$\tau_1 = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, X\} \text{ and } \tau_2 = \{\emptyset, \{1\}, \{1, 2\}, X\}.$$

Let $Y = \{a, b, c\}$ with topologies $\sigma_1 = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, Y\}$, $\sigma_2 = \{\emptyset, \{a\}, \{b\}, \{a, b\}, Y\}$. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is defined as follow: $F(1) = \{a, b\}$, $F(2) = \{a\}$, $F(3) = \{b\}$. Then, F is punctually (τ_1, τ_2) -paracompact.

Example 3.1.13. Let $X = \{1, 2, 3\}$ with topologies

$$\tau_1 = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, X\}$$

and $\tau_2 = \{\emptyset, \{1\}, \{1, 2\}, X\}$. Let $Y = \{a, b, c\}$ with topologies $\sigma_1 = \{\emptyset, \{b, c\}, Y\}$ and $\sigma_2 = \{\emptyset, \{a, c\}, Y\}$. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is defined as follow: $F(1) = F(2) = F(3) = \{a, b\}$. Then, F is punctually (τ_1, τ_2) -regular.

Lemma 3.1.14. Let (X, τ_1, τ_2) be a bitopological space. Then, $\tau_1\tau_2\text{-}\beta\text{Cl}(A) \subseteq \tau_1\tau_2\text{-Cl}(A)$ for every subset A of X .

Proof. Let $x \in X - \tau_1\tau_2\text{-Cl}(A)$. By Proposition 2.1.31, $x \in \tau_1\tau_2\text{-Int}(X - A)$ and there exists a $\tau_1\tau_2$ -open set V such that $x \in V \subseteq X - A$. Since every $\tau_1\tau_2$ -open set is $\tau_1\tau_2\text{-}\beta$ -open, we have $x \in \tau_1\tau_2\text{-}\beta\text{Int}(X - A)$. By Proposition 2.1.41(1), $x \in X - \tau_1\tau_2\text{-}\beta\text{Cl}(A)$, so $X - \tau_1\tau_2\text{-Cl}(A) \subseteq X - \tau_1\tau_2\text{-}\beta\text{Cl}(A)$. Consequently, we obtain $\tau_1\tau_2\text{-}\beta\text{Cl}(A) \subseteq \tau_1\tau_2\text{-Cl}(A)$. \square

Lemma 3.1.15. If $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is punctually (τ_1, τ_2) -paracompact and punctually (τ_1, τ_2) -regular, then $\beta\text{Cl}F_{*}^{+}(V) = F^{+}(V)$ for every $\sigma_1\sigma_2$ -open V of Y .

Proof. Let V be a $\sigma_1\sigma_2$ -open set of Y and $x \in \beta\text{Cl}F_{*}^{+}(V)$. Then, we have $\sigma_1\sigma_2\text{-}\beta\text{Cl}(F(x)) \subseteq V$ and $F(x) \subseteq V$. Therefore, we have $x \in F^{+}(V)$, and hence $\beta\text{Cl}F_{*}^{+}(V) \subseteq F^{+}(V)$. On the other hand, let $x \in F^{+}(V)$. Then, $F(x) \subseteq V$ and by Lemma 3.1.10, there exists a $\sigma_1\sigma_2$ -open set U of Y such that $F(x) \subseteq \sigma_1\sigma_2\text{-Cl}(U) \subseteq U \subseteq V$. By Lemma 3.1.14, we have $\sigma_1\sigma_2\text{-}\beta\text{Cl}(F(x)) \subseteq \sigma_1\sigma_2\text{-Cl}(U) \subseteq V$. This shows that $x \in \beta\text{Cl}F_{*}^{+}(V)$, and hence $F^{+}(V) \subseteq \beta\text{Cl}F_{*}^{+}(V)$. Consequently, we obtain $\beta\text{Cl}F_{*}^{+}(V) = F^{+}(V)$. \square

Theorem 3.1.16. Let $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ be punctually (τ_1, τ_2) -paracompact and punctually (τ_1, τ_2) -regular. Then, F is upper $\beta(\tau_1, \tau_2)$ -continuous if and only if $\beta\text{Cl}F_{*} : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is upper $\beta(\tau_1, \tau_2)$ -continuous.

Proof. Suppose that F is upper $\beta(\tau_1, \tau_2)$ -continuous. Let $x \in X$ and V be a $\sigma_1\sigma_2$ -open set of Y such that $\beta\text{Cl}F_{*}(x) \subseteq V$. By Lemma 3.1.15, we have $x \in \beta\text{Cl}F_{*}^{+}(V) = F^{+}(V)$. Since F is upper $\beta(\tau_1, \tau_2)$ -continuous, there exists a $\tau_1\tau_2\text{-}\beta$ -open set U containing x such that $F(U) \subseteq V$. Since $F(z)$ is $\sigma_1\sigma_2$ -paracompact and $\sigma_1\sigma_2$ -regular for each $z \in U$, by Lemma 3.1.10 there exists a $\sigma_1\sigma_2$ -open set H such that $F(z) \subseteq H \subseteq \sigma_1\sigma_2\text{-Cl}(H) \subseteq V$. By Lemma 3.1.14, we have $\sigma_1\sigma_2\text{-}\beta\text{Cl}(F(z)) \subseteq \sigma_1\sigma_2\text{-Cl}(H) \subseteq V$ for each $z \in U$, and hence $\beta\text{Cl}F_{*}(U) \subseteq V$. This shows that $\beta\text{Cl}F_{*}$ is upper $\beta(\tau_1, \tau_2)$ -continuous.

Conversely, we suppose that $\beta\text{Cl}F_{*} : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is upper $\beta(\tau_1, \tau_2)$ -continuous. Let $x \in X$ and V be a $\sigma_1\sigma_2$ -open set of Y such that $F(x) \subseteq V$. By

Lemma 3.1.15, we have $x \in F^+(V) = \beta ClF_{\otimes}^+(V)$, and hence $\beta ClF_{\otimes}(x) \subseteq V$. Since βClF_{\otimes} is upper $\beta(\tau_1, \tau_2)$ -continuous, there exists a $\tau_1\tau_2$ - β -open set U of containing x such that $\beta ClF_{\otimes}(U) \subseteq V$; hence, $F(U) \subseteq V$. This shows that F is upper $\beta(\tau_1, \tau_2)$ -continuous. \square

Lemma 3.1.17. For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, it follows that for each $\sigma_1\sigma_2$ - β -open set V of Y $\beta ClF_{\otimes}^-(V) = F^-(V)$.

Proof. Suppose that V is a $\sigma_1\sigma_2$ - β -open set Y . Let $x \in \beta ClF_{\otimes}^-(V)$. Then, $\sigma_1\sigma_2$ - $\beta Cl(F(x)) \cap V \neq \emptyset$. Hence, $F(x) \cap V \neq \emptyset$. Therefore, we obtain $x \in F^-(V)$. This shows that $\beta ClF_{\otimes}^-(V) \subseteq F^-(V)$. On the other hand, let $x \in F^-(V)$. Then, we have $\emptyset \neq F(x) \cap V \subseteq \sigma_1\sigma_2$ - $\beta Cl(F(x)) \cap V$. Thus, $x \in \beta ClF_{\otimes}^-(V)$. This shows that $F^-(V) \subseteq \beta ClF_{\otimes}^-(V)$. Consequently, we obtain $\beta ClF_{\otimes}^-(V) = F^-(V)$. \square

Theorem 3.1.18. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower $\beta(\tau_1, \tau_2)$ -continuous if and only if $\beta ClF_{\otimes} : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower $\beta(\tau_1, \tau_2)$ -continuous.

Proof. By utilizing Lemma 3.1.17, this can be proved similarly to that of Theorem 3.1.16. \square

For a multifunction $F : X \rightarrow Y$, the graph multifunction $G_F : X \rightarrow X \times Y$ is defined as follows: $G_F(x) = \{x\} \times F(x)$ for every $x \in X$.

Lemma 3.1.19. [10] The following hold for a multifunction $F : X \rightarrow Y$:

- (i) $G_F^+(A \times B) = A \cap F^+(B)$,
- (ii) $G_F^-(A \times B) = A \cap F^-(B)$,

for any subsets $A \subseteq X$ and $B \subseteq Y$.

Lemma 3.1.20. Let (X, τ_1, τ_2) be a bitopological space. If A is $\tau_1\tau_2$ - β -open and B is $\tau_1\tau_2$ -open in X , then $A \cap B$ is $\tau_1\tau_2$ - β -open.

Proof. Suppose that A is $\tau_1\tau_2$ - β -open and B is $\tau_1\tau_2$ -open in X . Then, we have $A \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(\tau_1\text{-Cl}(A)))$ and $B = \tau_1\text{-Int}(B) = \tau_2\text{-Int}(B)$. By Proposition 2.1.6(1),

$$A \cap B \subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(\tau_1\text{-Cl}(A))) \cap B$$

$$\begin{aligned}
&\subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(\tau_1\text{-Cl}(A)) \cap B) \\
&= \tau_1\text{-Cl}(\tau_2\text{-Int}(\tau_1\text{-Cl}(A \cap B))) \\
&\subseteq \tau_1\text{-Cl}(\tau_2\text{-Int}(\tau_1\text{-Cl}(A \cap B))).
\end{aligned}$$

Consequently, we obtain $A \cap B$ is $\tau_1\tau_2$ - β -open. \square

Definition 3.1.21. [21] A bitopological space (X, τ_1, τ_2) is said to be $\tau_1\tau_2$ -compact if every cover of X by $\tau_1\tau_2$ -open sets of X has a finite subcover.

By ρ_i we denote the product topology $\tau_i \times \sigma_i$ for $i = 1, 2$.

Theorem 3.1.22. Let $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ be a multifunction such that $F(x)$ is $\sigma_1\sigma_2$ -compact for each $x \in X$. Then, F is upper $\beta(\tau_1, \tau_2)$ -continuous if and only if $G_F : (X, \tau_1, \tau_2) \rightarrow (X \times Y, \rho_1, \rho_2)$ is upper $\beta(\tau_1, \tau_2)$ -continuous.

Proof. Suppose that $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is upper $\beta(\tau_1, \tau_2)$ -continuous. Let $x \in X$ and W be a $\rho_1\rho_2$ -open set of $X \times Y$ containing $G_F(x)$. For each $y \in F(x)$, there exist $\tau_1\tau_2$ -open set $U(y)$ of X and $\sigma_1\sigma_2$ -open set $V(y)$ of Y such that $(x, y) \in U(y) \times V(y) \subseteq W$. The family $\{V(y) \mid y \in F(x)\}$ is $\sigma_1\sigma_2$ -open cover of $F(x)$ and there exists a finite number of points, say, y_1, y_2, \dots, y_n in $F(x)$ such that $F(x) \subseteq \cup\{V(y_i) \mid 1 \leq i \leq n\}$. Put

$$U = \cap\{U(y_i) \mid 1 \leq i \leq n\} \text{ and } V = \cup\{V(y_i) \mid 1 \leq i \leq n\}.$$

Then, we have U is $\tau_1\tau_2$ -open in X and V is $\sigma_1\sigma_2$ -open in Y such that $\{x\} \times F(x) \subseteq U \times V \subseteq W$. Since F is upper $\beta(\tau_1, \tau_2)$ -continuous, there exists a $\tau_1\tau_2$ - β -open set G containing x such that $F(G) \subseteq V$. By Lemma 3.1.19 and 2.2.4(3), we have $U \cap G \subseteq U \cap F^+(V) = G_F^+(U \times V) \subseteq G_F^+(W)$. By Lemma 3.1.20, $U \cap G$ is $\tau_1\tau_2$ - β -open in X , and $G_F(U \cap G) \subseteq W$ by Lemma 2.2.4(1). This shows that G_F is upper $\beta(\tau_1, \tau_2)$ -continuous.

Conversely, we suppose that $G_F : (X, \tau_1, \tau_2) \rightarrow (X \times Y, \rho_1, \rho_2)$ is upper $\beta(\tau_1, \tau_2)$ -continuous. Let $x \in X$ and V be a $\sigma_1\sigma_2$ -open containing $F(x)$. Since $X \times V$ is $\rho_1\rho_2$ -open in $X \times Y$ and $G_F(x) \subseteq X \times V$, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $G_F(U) \subseteq X \times V$. Therefore, by Lemma 2.2.4(1) and 3.1.19, $U \subseteq G_F^+(X \times V) = F^+(V)$, and $F(U) \subseteq V$. This shows that F is upper $\beta(\tau_1, \tau_2)$ -continuous. \square

Theorem 3.1.23. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower $\beta(\tau_1, \tau_2)$ -continuous if and only if $G_F : (X, \tau_1, \tau_2) \rightarrow (X \times Y, \rho_1, \rho_2)$ is lower $\beta(\tau_1, \tau_2)$ -continuous.

Proof. Suppose that $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower $\beta(\tau_1, \tau_2)$ -continuous. Let $x \in X$ and W be a $\rho_1\rho_2$ -open set of $X \times Y$ such that $G_F(x) \cap W \neq \emptyset$. There exists $y \in F(x)$ such that $(x, y) \in W$, and hence $(x, y) \in U \times V \subseteq W$ for some $\tau_1\tau_2$ -open set U of X and $\sigma_1\sigma_2$ -open set V of Y . Since $F(x) \cap V \neq \emptyset$, there exists a $\tau_1\tau_2$ - β -open set G containing x such that $F(z) \cap V \neq \emptyset$ for each $z \in G$; hence $G \subseteq F^-(V)$. By Lemma 3.1.19 and Lemma 3.1.20, we have $U \cap G \subseteq U \cap F^-(V) = G_F^-(U \times V) \subseteq G_F^-(W)$. Moreover, $U \cap G$ is a $\tau_1\tau_2$ - β -open set containing x , and hence G_F is lower $\beta(\tau_1, \tau_2)$ -continuous.

Conversely, we suppose that $G_F : (X, \tau_1, \tau_2) \rightarrow (X \times Y, \rho_1, \rho_2)$ is lower $\beta(\tau_1, \tau_2)$ -continuous. Let $x \in X$ and V be a $\sigma_1\sigma_2$ -open set of Y such that $F(x) \cap V \neq \emptyset$. Then, we have $X \times V$ is $\rho_1\rho_2$ -open in $X \times Y$ and

$$G_F(x) \cap (X \times V) = (\{x\} \times F(x)) \cap (X \times V) = \{x\} \times (F(x) \cap V) \neq \emptyset.$$

There exists a $\tau_1\tau_2$ - β -open set U containing x such that $G_F(z) \cap (X \times V) \neq \emptyset$ for each $z \in U$. By Lemma 3.1.19, we obtain $U \subseteq G_F^-(X \times V) = F^-(V)$.

This shows that F is lower $\beta(\tau_1, \tau_2)$ -continuous. \square

3.2 Characterizations of upper and lower almost $\beta(\tau_1, \tau_2)$ -continuous multifunctions

In this section, we introduce the concepts of upper and lower almost $\beta(\tau_1, \tau_2)$ -continuous multifunctions. Moreover, several interesting characterizations of these multifunctions are discussed.

Definition 3.2.1. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is said to be:

- (1) *upper almost $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(U) \subseteq \sigma_1\text{-Int}(\sigma_2\text{-Cl}(V))$;
- (2) *lower almost $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V

of Y such that $F(x) \cap V \neq \emptyset$, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(z) \cap \sigma_1\text{-Int}(\sigma_2\text{-Cl}(V)) \neq \emptyset$ for every $z \in U$;

- (3) *upper almost (resp. lower almost) $\beta(\tau_1, \tau_2)$ -continuous* if F has this property at each point of X .

Remark. For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following implication holds:

$$\text{upper } \beta(\tau_1, \tau_2)\text{-continuity} \Rightarrow \text{upper almost } \beta(\tau_1, \tau_2)\text{-continuity}.$$

The converse of the implication is not true in general. We give an example for the implication as follows.

Example 3.2.2. Let $X = \{a, b, c\}$ with topologies $\tau_1 = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$ and $\tau_2 = \{\emptyset, \{a\}, \{a, b\}, X\}$. Let $Y = \{1, 2, 3, 4, 5\}$ with topologies $\sigma_1 = \{\emptyset, \{1\}, \{2, 3, 4, 5\}, Y\}$ and $\sigma_2 = \{\emptyset, \{2\}, \{3\}, \{2, 3\}, \{2, 3, 4, 5\}, Y\}$.

A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is defined as follow:

$F(a) = \{1\}$, $F(b) = \{2, 3\}$, $F(c) = \{4, 5\}$. Then, F is upper almost $\beta(\tau_1, \tau_2)$ -continuous, but F is not upper $\beta(\tau_1, \tau_2)$ -continuous.

Remark. For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following implication holds:

$$\text{lower } \beta(\tau_1, \tau_2)\text{-continuity} \Rightarrow \text{lower almost } \beta(\tau_1, \tau_2)\text{-continuity}.$$

The converse of the implication is not true in general. We give an example for the implication as follows.

Example 3.2.3. Let $X = \{a, b\}$ with topologies $\tau_1 = \{\emptyset, \{b\}, X\}$ and $\tau_2 = \{\emptyset, X\}$. Let $Y = \{1, 2, 3, 4\}$ with topologies $\sigma_1 = \{\emptyset, \{1, 2\}, Y\}$ and $\sigma_2 = \{\emptyset, \{1\}, \{1, 2\}, Y\}$. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is defined as follow: $F(a) = \{1, 2\}$, $F(b) = \{3\}$. Then, F is lower almost $\beta(\tau_1, \tau_2)$ -continuous, but F is not lower $\beta(\tau_1, \tau_2)$ -continuous.

Theorem 3.2.4. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is upper almost $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-sCl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$.

Proof. Let V be a $\sigma_1\sigma_2$ -open set containing $F(x)$. Then, there exists a $\tau_1\tau_2\text{-}\beta$ -open set U containing x such that $F(U) \subseteq \sigma_1\sigma_2\text{-sCl}(V)$. Then, $x \in U \subseteq F^+(\sigma_1\sigma_2\text{-sCl}(V))$. Therefore, $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-sCl}(V)))$.

Conversely, let V be a $\sigma_1\sigma_2$ -open set containing $F(x)$. Moreover, we have $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-sCl}(V)))$. There exists a $\tau_1\tau_2\text{-}\beta$ -open set G containing x such that $G \subseteq F^+(\sigma_1\sigma_2\text{-sCl}(V))$, and hence $F(G) \subseteq \sigma_1\sigma_2\text{-sCl}(V) = \sigma_1\text{-Int}(\sigma_2\text{-Cl}(V))$. This shows that F is upper almost $\beta(\tau_1, \tau_2)$ -continuous at x . \square

Theorem 3.2.5. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower almost $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^-(\sigma_1\sigma_2\text{-sCl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y such that $F(x) \cap V \neq \emptyset$.

Proof. The proof is similar to that of Theorem 3.2.4. \square

Theorem 3.2.6. For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is upper almost $\beta(\tau_1, \tau_2)$ -continuous;
- (2) for each $x \in X$ and each $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$, there exists a $\tau_1\tau_2\text{-}\beta$ -open set U of X containing x such that $F(U) \subseteq \sigma_1\sigma_2\text{-sCl}(V)$;
- (3) $F^+(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-sCl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y ;
- (4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-sInt}(K))) \subseteq F^-(K)$ for every $\sigma_1\sigma_2$ -closed set K of Y .

Proof. (1) \Rightarrow (2): The proof follows from Definition 3.2.1(1).

(2) \Rightarrow (3): Let V be a $\sigma_1\sigma_2$ -open set of Y and $x \in F^+(V)$. Then, $F(x) \subseteq V$ and there exists a $\tau_1\tau_2\text{-}\beta$ -open set U containing x such that $F(U) \subseteq \sigma_1\text{-Int}(\sigma_2\text{-Cl}(V)) = \sigma_1\sigma_2\text{-sCl}(V)$. Therefore, we have $x \in U \subseteq F^+(\sigma_1\sigma_2\text{-sCl}(V))$. Thus,

$$x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-sCl}(V))).$$

Consequently, we obtain $F^+(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-sCl}(V)))$.

(3) \Rightarrow (4): Let K be a $\sigma_1\sigma_2$ -closed set of Y . Since $Y - K$ is $\sigma_1\sigma_2$ -open, by Lemma 2.2.4(5) we obtain

$$\begin{aligned} X - F^-(K) &= F^+(Y - K) \\ &\subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-sCl}(Y - K))) \\ &= \tau_1\tau_2\text{-}\beta\text{Int}(F^+(Y - \sigma_1\sigma_2\text{-sInt}(K))) \\ &= \tau_1\tau_2\text{-}\beta\text{Int}(X - F^-(\sigma_1\sigma_2\text{-sInt}(K))) \\ &= X - \tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-sInt}(K))). \end{aligned}$$

Therefore, we obtain $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-sInt}(K))) \subseteq F^-(K)$.

(4) \Rightarrow (3): The proof is obvious.

(3) \Rightarrow (1): Let $x \in X$ and V be a $\sigma_1\sigma_2$ -open set containing $F(x)$. Then, we have $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-sCl}(V)))$. Therefore, by Theorem 3.2.4, F is upper almost $\beta(\tau_1, \tau_2)$ -continuous at x . \square

Theorem 3.2.7. For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is lower almost $\beta(\tau_1, \tau_2)$ -continuous;
- (2) for each $x \in X$ and each $\sigma_1\sigma_2$ -open set V containing $F(x)$, there exists a $\tau_1\tau_2\text{-}\beta$ -open set U containing x such that $F(z) \cap \sigma_1\sigma_2\text{-sCl}(V)$ for each $z \in U$;
- (3) $F^-(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^-(\sigma_1\sigma_2\text{-sCl}(V)))$ for every $\sigma_1\sigma_2$ -open set V ;
- (4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-sInt}(K))) \subseteq F^+(K)$ for every $\sigma_1\sigma_2$ -closed set K .

Proof. By utilizing Theorem 3.2.5, this can be similar to Theorem 3.2.6. \square

Theorem 3.2.8. If a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower almost $\beta(\tau_1, \tau_2)$ -continuous, then $\beta\text{Cl}F_{\otimes} : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower almost $\beta(\tau_1, \tau_2)$ -continuous.

Proof. Suppose that F is lower almost $\beta(\tau_1, \tau_2)$ -continuous. Let $x \in X$ and V be a $\sigma_1\sigma_2$ -open set of Y such that $\beta\text{Cl}F_{\otimes}(x) \cap V \neq \emptyset$. By Lemma 3.1.17, we have

$x \in \beta \text{Cl} F_{\otimes}^{-}(V) = F^{-}(V)$. Since F is lower almost $\beta(\tau_1, \tau_2)$ -continuous, there exists a $\tau_1 \tau_2$ - β -open set U containing x such that

$$F(z) \cap \sigma_1\text{-Int}(\sigma_2\text{-Cl}(V)) \neq \emptyset \text{ for each } z \in U.$$

Therefore, $\sigma_1 \sigma_2$ - $\beta \text{Cl}(F(z)) \cap \sigma_1\text{-Int}(\sigma_2\text{-Cl}(V)) \neq \emptyset$ for each $z \in U$, and hence

$$\beta \text{Cl} F_{\otimes}(z) \cap \sigma_1\text{-Int}(\sigma_2\text{-Cl}(V)) \neq \emptyset \text{ for each } z \in U.$$

This shows that $\beta \text{Cl} F_{\otimes}$ is lower almost $\beta(\tau_1, \tau_2)$ -continuous. \square

Definition 3.2.9. Let (X, τ_1, τ_2) be a bitopological space. The β -frontier of a subset A of X , denoted by $\tau_1 \tau_2$ - $\beta \text{Fr}(A)$, is defined by

$$\begin{aligned} \tau_1 \tau_2\text{-}\beta \text{Fr}(A) &= \tau_1 \tau_2\text{-}\beta \text{Cl}(A) \cap \tau_1 \tau_2\text{-}\beta \text{Cl}(X - A) \\ &= \tau_1 \tau_2\text{-}\beta \text{Cl}(A) - \tau_1 \tau_2\text{-}\beta \text{Int}(A). \end{aligned}$$

Theorem 3.2.10. The set of all points x of X at which a multifunction

$F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is not upper $\beta(\tau_1, \tau_2)$ -continuous is identical with the union of the $\tau_1 \tau_2$ - β -frontier of the upper inverse images of $\sigma_1 \sigma_2$ -open sets containing $F(x)$.

Proof. Let $x \in X$ at which F is not upper $\beta(\tau_1, \tau_2)$ -continuous. There exists a $\sigma_1 \sigma_2$ -open set V containing $F(x)$ such that $U \cap (X - F^+(V)) \neq \emptyset$ for every $\tau_1 \tau_2$ - β -open set U containing x . Then, we have

$$x \in \tau_1 \tau_2\text{-}\beta \text{Cl}(X - F^+(V)) = X - \tau_1 \tau_2\text{-}\beta \text{Int}(F^+(V))$$

and $x \in F^+(V)$. Hence, we obtain $x \in \tau_1 \tau_2\text{-}\beta \text{Fr}(F^+(V))$.


Conversely, we suppose that V is a $\sigma_1 \sigma_2$ -open set containing $F(x)$ such that $x \in \tau_1 \tau_2\text{-}\beta \text{Fr}(F^+(V))$. If F is upper $\beta(\tau_1, \tau_2)$ -continuous at x , there exists a $\tau_1 \tau_2$ - β -open set U containing x such that $U \subseteq F^+(V)$. This implies that $x \in \tau_1 \tau_2\text{-}\beta \text{Int}(F^+(V))$. This is a contradiction; hence, F is not upper $\beta(\tau_1, \tau_2)$ -continuous. \square

Theorem 3.2.11. The set of all points x of X at which a multifunction

$F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is not lower $\beta(\tau_1, \tau_2)$ -continuous is identical with the

union of the $\tau_1\tau_2$ - β -frontier of the lower inverse images of $\sigma_1\sigma_2$ -open sets meeting $F(x)$.

Proof. The proof is similar to that of Theorem 3.2.10. □



CHAPTER 4

WEAKLY $\beta(\tau_1, \tau_2)$ -CONTINUOUS MULTIFUNCTIONS

4.1 Weakly $\beta(\tau_1, \tau_2)$ -continuous multifunctions

In this section, we introduce and investigate the notions of upper and lower weakly $\beta(\tau_1, \tau_2)$ -continuous multifunctions. Furthermore, the relationships weak $\beta(\tau_1, \tau_2)$ -continuity and the other types of continuity are investigated.

Definition 4.1.1. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is said to be:

- (1) *upper weakly $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(U) \subseteq \sigma_1\sigma_2\text{-Cl}(V)$;
- (2) *lower weakly $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y such that $F(x) \cap V \neq \emptyset$, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(z) \cap \sigma_1\sigma_2\text{-Cl}(V) \neq \emptyset$ for every $z \in U$;
- (3) *upper weakly (resp. lower weakly) $\beta(\tau_1, \tau_2)$ -continuous* if F has this property at each point of X .

Theorem 4.1.2. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is upper weakly $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$.

Proof. Let V be a $\sigma_1\sigma_2$ -open set containing $F(x)$. Therefore, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(U) \subseteq \sigma_1\sigma_2\text{-Cl}(V)$. Therefore, $x \in U \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(V))$. Since U is $\tau_1\tau_2$ - β -open, we have $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-Cl}(V)))$. Conversely, let V be a $\sigma_1\sigma_2$ -open set of Y containing $F(x)$, and we have $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-Cl}(V)))$. There exists a $\tau_1\tau_2$ - β -open set G of X containing x such that $G \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(V))$; hence, $F(G) \subseteq \sigma_1\sigma_2\text{-Cl}(V)$. This shows that F is upper weakly $\beta(\tau_1, \tau_2)$ -continuous at x . \square

Theorem 4.1.3. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower weakly $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^-(\sigma_1\sigma_2\text{Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y such that $F(x) \cap V \neq \emptyset$.

Proof. The proof is similar to that of Theorem 4.1.2. □

Remark. For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following implication hold:

$$\begin{array}{c} \text{upper } (\tau_1, \tau_2)\text{-}\beta\text{-continuity} \\ \Downarrow \\ \text{upper almost } (\tau_1, \tau_2)\text{-}\beta\text{-continuity} \\ \Downarrow \\ \text{upper weak } (\tau_1, \tau_2)\text{-}\beta\text{-continuity.} \end{array}$$

The converse of the implications are not true in general. We give an example for the implication as follows.

Example 4.1.4. Let $X = \{1, 2, 3\}$ with topologies $\tau_1 = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, X\}$ and $\tau_2 = \{\emptyset, \{2, 3\}, X\}$. Let $Y = \{a, b, c\}$ with topologies $\sigma_1 = \{\emptyset, \{a\}, \{c\}, \{a, c\}, Y\}$ and $\sigma_2 = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, c\}, \{a, b\}, \{b, c\}, Y\}$. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is defined as follow: $F(1) = \{b\}$, $F(2) = \{c\}$, $F(3) = \{a, b\}$. Then, F is upper(lower) weakly $\beta(\tau_1, \tau_2)$ -continuous, but F is not upper almost $\beta(\tau_1, \tau_2)$ -continuous and upper $\beta(\tau_1, \tau_2)$ -continuous.

The following theorems give some characterizations of upper and lower weakly $\beta(\tau_1, \tau_2)$ -continuous multifunctions.

Theorem 4.1.5. For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is upper weakly $\beta(\tau_1, \tau_2)$ -continuous;
- (2) $F^+(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y ;
- (3) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(K))) \subseteq F^-(K)$ for every $\sigma_1\sigma_2$ -closed set K of Y ;

(4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(B)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B of Y ;

(5) $F^+(\sigma_1\sigma_2\text{-Int}(B)) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-Cl}(\sigma_1\sigma_2\text{-Int}(B))))$ for every subset B of Y ;

(6) $F^+(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y ;

(7) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(V)) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(V))$ for every $\sigma_1\sigma_2$ -open set V of Y .

Proof. (1) \Rightarrow (2): Let V be a $\sigma_1\sigma_2$ -open set of Y and $x \in F^+(V)$. Then, $F(x) \subseteq V$ and by Theorem 4.1.2, $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{Cl}(V)))$. Therefore, we obtain $F^+(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{Cl}(V)))$.

(2) \Rightarrow (3): Let K be a $\sigma_1\sigma_2$ -closed set of Y . Therefore, $Y - K$ is a $\sigma_1\sigma_2$ -open set. By (2),

$$\begin{aligned} X - F^-(K) &= F^+(Y - K) \\ &\subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{Cl}(Y - K))) \\ &= X - \tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{Int}(K))). \end{aligned}$$

Consequently, we obtain $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{Int}(K))) \subseteq F^-(K)$.

(3) \Rightarrow (4): Let B be a subset of Y . Then, $\sigma_1\sigma_2\text{-Cl}(B)$ is $\sigma_1\sigma_2$ -closed in Y . Thus, we obtain $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(B)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(B))$.

(4) \Rightarrow (5): Let B be a subset of Y . By (4), we obtain

$$\begin{aligned} F^+(\sigma_1\sigma_2\text{-Int}(B)) &= X - F^-(\sigma_1\sigma_2\text{-Cl}(Y - B)) \\ &\subseteq X - \tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(Y - B)))) \\ &= \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-Cl}(\sigma_1\sigma_2\text{-Int}(B)))). \end{aligned}$$

(5) \Rightarrow (6): The proof is obvious.

(6) \Rightarrow (7): Let V be a $\sigma_1\sigma_2$ -open set of Y . By (6), we have

$$\begin{aligned} \tau_1\tau_2\text{-}\beta\text{Cl}(F^-(V)) &\subseteq \tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \\ &= \tau_1\tau_2\text{-}\beta\text{Cl}(X - F^+(Y - \sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \end{aligned}$$

$$\begin{aligned}
&= X - \tau_1\tau_2\text{-}\beta\text{Int}(F^+(Y - \sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \\
&= X - \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-Cl}(Y - \sigma_1\sigma_2\text{-Cl}(V)))) \\
&\subseteq X - F^+(Y - \sigma_1\sigma_2\text{-Cl}(V)) \\
&= F^-(\sigma_1\sigma_2\text{-Cl}(V)).
\end{aligned}$$

Consequently, we obtain $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(V)) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(V))$.

(7) \Rightarrow (1): Let $x \in X$ and V be a $\sigma_1\sigma_2$ -open set containing $F(x)$. By (7), we have

$$\begin{aligned}
x \in F^+(V) &\subseteq F^+(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V))) \\
&= X - F^-(\sigma_1\sigma_2\text{-Cl}(Y - \sigma_1\sigma_2\text{-Cl}(V))) \\
&\subseteq X - \tau_1\tau_2\text{-}\beta\text{Cl}(F^-(Y - \sigma_1\sigma_2\text{-Cl}(V))) \\
&= \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-Cl}(V))).
\end{aligned}$$

Therefore, $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-Cl}(V)))$, and hence F is upper weakly $\beta(\tau_1, \tau_2)$ -continuous by Theorem 4.1.2. \square

Theorem 4.1.6. For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is lower weakly $\beta(\tau_1, \tau_2)$ -continuous;
- (2) $F^-(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^-(\sigma_1\sigma_2\text{-Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y ;
- (3) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-Int}(K)))) \subseteq F^+(K)$ for every $\sigma_1\sigma_2$ -closed set K of Y ;
- (4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(B)))) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B of Y ;
- (5) $F^-(\sigma_1\sigma_2\text{-Int}(B)) \subseteq \tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Cl}(\sigma_1\sigma_2\text{-Int}(B))))$ for every subset B of Y ;
- (6) $F^-(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^-(\sigma_1\sigma_2\text{-Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y ;
- (7) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(V)) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(V))$ for every $\sigma_1\sigma_2$ -open set V of Y .

Proof. The proof is similar to that of Theorem 4.1.5. \square

Definition 4.1.7. Let A be a subset of a bitopological space (X, τ_1, τ_2) . A point $x \in X$ is called $\tau_1\tau_2$ - θ -cluster point of A if $\tau_1\tau_2\text{-Cl}(U) \cap A \neq \emptyset$ for every $\tau_1\tau_2$ -open set U containing x . The set of all $\tau_1\tau_2$ - θ -cluster point of A is called $\tau_1\tau_2$ - θ -closure of A and is denoted by $\tau_1\tau_2\text{-Cl}_\theta(A)$.

A subset A of X is said to be $\tau_1\tau_2$ - θ -closed if $A = \tau_1\tau_2\text{-Cl}_\theta(A)$. The complement of a $\tau_1\tau_2$ - θ -closed set is said to be $\tau_1\tau_2$ - θ -open. The union of all $\tau_1\tau_2$ - θ -open sets contained in A is called $\tau_1\tau_2$ - θ -interior of A and is denoted by $\tau_1\tau_2\text{-Int}_\theta(A)$.

Lemma 4.1.8. For a subset A of a bitopological space (X, τ_1, τ_2) , the following properties hold:

- (1) If A is $\tau_1\tau_2$ -open in X , then $\tau_1\tau_2\text{-Cl}(A) = \tau_1\tau_2\text{-Cl}_\theta(A)$.
- (2) $\tau_1\tau_2\text{-Cl}_\theta(A)$ is $\tau_1\tau_2$ -closed in X .

Proof. (1) In general, this holds that $\tau_1\tau_2\text{-Cl}(A) \subseteq \tau_1\tau_2\text{-Cl}_\theta(A)$. Suppose that $x \notin \tau_1\tau_2\text{-Cl}(A)$. Then, there exists a $\tau_1\tau_2$ -open set U containing x such that $U \cap A = \emptyset$; hence $\tau_1\tau_2\text{-Cl}(U) \cap A = \emptyset$. This shows that $x \notin \tau_1\tau_2\text{-Cl}_\theta(A)$. Therefore, we obtain $\tau_1\tau_2\text{-Cl}_\theta(A) \subseteq \tau_1\tau_2\text{-Cl}(A)$. Consequently, $\tau_1\tau_2\text{-Cl}(A) = \tau_1\tau_2\text{-Cl}_\theta(A)$.

(2) Let $x \in X - \tau_1\tau_2\text{-Cl}_\theta(A)$. Then, $x \notin \tau_1\tau_2\text{-Cl}_\theta(A)$. There exists a $\tau_1\tau_2$ -open set U_x containing x such that $\tau_1\tau_2\text{-Cl}(U_x) \cap A = \emptyset$. Then, we have $\tau_1\tau_2\text{-Cl}_\theta(A) \cap U_x = \emptyset$ and so $x \in U_x \subseteq X - \tau_1\tau_2\text{-Cl}_\theta(A)$. Therefore, we obtain $X - \tau_1\tau_2\text{-Cl}_\theta(A) = \bigcup_{x \in X - \tau_1\tau_2\text{-Cl}_\theta(A)} U_x$. This shows that $\tau_1\tau_2\text{-Cl}_\theta(A)$ is $\tau_1\tau_2$ -closed. \square

Definition 4.1.9. A subset A of a bitopological space (X, τ_1, τ_2) is said to be (τ_1, τ_2) - r -closed (resp. (τ_1, τ_2) - s -open, (τ_1, τ_2) - p -open, (τ_1, τ_2) - β -open) if $A = \tau_1\tau_2\text{-Cl}(\tau_1\tau_2\text{-Int}(A))$ (resp. $A \subseteq \tau_1\tau_2\text{-Cl}(\tau_1\tau_2\text{-Int}(A))$, $A \subseteq \tau_1\tau_2\text{-Int}(\tau_1\tau_2\text{-Cl}(A))$, $A \subseteq \tau_1\tau_2\text{-Cl}(\tau_1\tau_2\text{-Int}(\tau_1\tau_2\text{-Cl}(A)))$)

Theorem 4.1.10. For a multifunction $F: (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is upper weakly $\beta(\tau_1, \tau_2)$ -continuous;
- (2) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}_\theta(B)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}_\theta(B))$ for every subset B of Y ;

- (3) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(B)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}_\theta(B))$ for every subset B of Y ;
- (4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(V))$ for every $\sigma_1\sigma_2$ -open set V of Y ;
- (5) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(V))$ for every (σ_1, σ_2) p-open set V of Y ;
- (6) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(K))) \subseteq F^-(K)$ for every (σ_1, σ_2) r-closed set K of Y ;
- (7) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(V))$ for every $(\sigma_1, \sigma_2)\beta$ -open set V of Y ;
- (8) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(V))$ for every (σ_1, σ_2) s-open set V of Y .

Proof. (1) \Rightarrow (2): Let B be any subset of Y . Then, $\sigma_1\sigma_2\text{-Cl}_\theta(B)$ is $\sigma_1\sigma_2$ -closed in Y . Therefore, by Theorem 4.1.5(3) we obtain

$$\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}_\theta(B)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}_\theta(B)).$$

(2) \Rightarrow (3): This is obvious since $\sigma_1\sigma_2\text{-Cl}(B) \subseteq \sigma_1\sigma_2\text{-Cl}_\theta(B)$ for every subset B of Y .

(3) \Rightarrow (4): This is obvious since $\sigma_1\sigma_2\text{-Cl}(V) = \sigma_1\sigma_2\text{-Cl}_\theta(V)$ for every $\sigma_1\sigma_2$ -open set V of Y .

(4) \Rightarrow (5): Let V be a (σ_1, σ_2) p-open set of Y . Then, we have $V \subseteq \sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V))$, and hence

$$\sigma_1\sigma_2\text{-Cl}(V) = \sigma_1\sigma_2\text{-Cl}(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V))).$$

Now, put $G = \sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V))$, then G is $\sigma_1\sigma_2$ -open in Y and $\sigma_1\sigma_2\text{-Cl}(G) = \sigma_1\sigma_2\text{-Cl}(V)$. Therefore, by (4) we have

$$\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(V)).$$

(5) \Rightarrow (6): Let K be any (σ_1, σ_2) r-closed set of Y . Then, we have $\sigma_1\sigma_2\text{-Int}(K)$ is (σ_1, σ_2) p-open in Y and by (5), we obtain

$$\begin{aligned}\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(K))) &= \tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(\sigma_1\sigma_2\text{-Int}(K))))) \\ &\subseteq F^-(\sigma_1\sigma_2\text{-Cl}(\sigma_1\sigma_2\text{-Int}(K))) = F^-(K).\end{aligned}$$

(6) \Rightarrow (7): Let V be any $(\sigma_1, \sigma_2)\beta$ -open set of Y . Then, we have

$$V \subseteq \sigma_1\sigma_2\text{-Cl}(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V))).$$

Since $\sigma_1\sigma_2\text{-Cl}(V)$ is (σ_1, σ_2) r-closed in Y , by (6)

$$\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(V)).$$

(7) \Rightarrow (8): This is obvious since every (σ_1, σ_2) s-open set is $(\sigma_1, \sigma_2)\beta$ -open.

(8) \Rightarrow (1): Let V be any $\sigma_1\sigma_2$ -open set of Y . Since V is (σ_1, σ_2) s-open set in Y , by (8) we have

$$\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(V)) \subseteq \tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(V)).$$


By Theorem 4.1.5, we obtain F is upper weakly $(\tau_1, \tau_2)\text{-}\beta$ -continuous. \square

Theorem 4.1.11. For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is lower weakly $\beta(\tau_1, \tau_2)$ -continuous;
- (2) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}_\theta(B)))) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}_\theta(B))$ for every subset B of Y ;
- (3) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(B)))) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}_\theta(B))$ for every subset B of Y ;
- (4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(V))$ for every $\sigma_1\sigma_2$ -open set V of Y ;
- (5) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(V))$ for every (σ_1, σ_2) p-open set V of Y ;

- (6) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-Int}(K))) \subseteq F^+(K)$ for every (σ_1, σ_2) r-closed set K of Y ;
- (7) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(V))$ for every (σ_1, σ_2) β -open set V of Y ;
- (8) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(V)))) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(V))$ for every (σ_1, σ_2) s-open set V of Y .

Proof. The proof is similar to that of Theorem 4.1.10. □



CHAPTER 5

CONCLUSIONS

5.1 Conclusions

The purposes of this thesis are to introduce the notions of $\beta(\tau_1, \tau_2)$ -continuous multifunctions, almost $\beta(\tau_1, \tau_2)$ -continuous multifunctions, and weakly $\beta(\tau_1, \tau_2)$ -continuous multifunctions; moreover, some characterizations of these multifunctions are obtained. The results are as follows:

1. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is said to be:

(1) *upper $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(U) \subseteq V$;

(2) *lower $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y such that $F(x) \cap V \neq \emptyset$, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(z) \cap V \neq \emptyset$ for every $z \in U$;

(3) *upper (resp. lower) $\beta(\tau_1, \tau_2)$ -continuous* if F has this property at each point of X .

From the above definition, the following theorems are derived:

1.1 A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is upper $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2$ - $\beta\text{Int}(F^+(V))$ for every $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$.

1.2 A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2$ - $\beta\text{Int}(F^-(V))$ for every $\sigma_1\sigma_2$ -open set V of Y such that $F(x) \cap V \neq \emptyset$.

1.3 For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is upper $\beta(\tau_1, \tau_2)$ -continuous;
- (2) $F^+(V)$ is $\tau_1\tau_2$ - β -open in X for every $\sigma_1\sigma_2$ -open set V of Y ;
- (3) $F^-(K)$ is $\tau_1\tau_2$ - β -closed in X for every $\sigma_1\sigma_2$ -closed set K of Y ;
- (4) $\tau_1\tau_2$ - $\beta\text{Cl}(F^-(B)) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B of Y ;

(5) $\tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\text{-Int}(F^-(B)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B .

1.4 For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is lower $\beta(\tau_1, \tau_2)$ -continuous;
- (2) $F^-(V)$ is $\tau_1\tau_2$ - β -open in X for every $\sigma_1\sigma_2$ -open set V of Y ;
- (3) $F^+(K)$ is $\tau_1\tau_2$ - β -closed in X for every $\sigma_1\sigma_2$ -closed set K of Y ;
- (4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(B)) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B of Y ;
- (5) $\tau_1\text{-Int}(\tau_2\text{-Cl}(\tau_1\text{-Int}(F^+(B)))) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B .

2. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is said to be:

- (1) *upper almost $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(U) \subseteq \sigma_1\text{-Int}(\sigma_2\text{-Cl}(V))$;
- (2) *lower almost $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y such that $F(x) \cap V \neq \emptyset$, there exists a $\tau_1\tau_2$ - β -open set U containing x such that $F(z) \cap \sigma_1\text{-Int}(\sigma_2\text{-Cl}(V)) \neq \emptyset$ for every $z \in U$;
- (3) *upper almost (resp. lower almost) $\beta(\tau_1, \tau_2)$ -continuous* if F has this property at each point of X .

From the above definition, the following theorems are derived:

2.1 A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is upper almost $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-sCl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$.

2.2 A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower almost $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^-(\sigma_1\sigma_2\text{-sCl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y such that $F(x) \cap V \neq \emptyset$.

2.3 For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is upper almost $\beta(\tau_1, \tau_2)$ -continuous;
- (2) for each $x \in X$ and each $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$, there exists a $\tau_1\tau_2$ - β -open set U of X containing x such that $F(U) \subseteq \sigma_1\sigma_2\text{-sCl}(V)$;
- (3) $F^+(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-sCl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y ;

(4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-sInt}(K))) \subseteq F^-(K)$ for every $\sigma_1\sigma_2$ -closed set K .

2.4 For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is lower almost $\beta(\tau_1, \tau_2)$ -continuous;
- (2) for each $x \in X$ and each $\sigma_1\sigma_2$ -open set V containing $F(x)$, there exists a $\tau_1\tau_2\text{-}\beta$ -open set U containing x such that $F(z) \cap \sigma_1\sigma_2\text{-sCl}(V)$ for each $z \in U$;
- (3) $F^-(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^-(\sigma_1\sigma_2\text{-sCl}(V)))$ for every $\sigma_1\sigma_2$ -open set V ;
- (4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-sInt}(K))) \subseteq F^+(K)$ for every $\sigma_1\sigma_2$ -closed set K .

2.5 If a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower almost $\beta(\tau_1, \tau_2)$ -continuous, then $\beta\text{Cl}F_{\otimes} : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower almost $\beta(\tau_1, \tau_2)$ -continuous.

3. A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is said to be:

- (1) *upper weakly $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$, there exists a $\tau_1\tau_2\text{-}\beta$ -open set U containing x such that $F(U) \subseteq \sigma_1\sigma_2\text{-Cl}(V)$;
- (2) *lower weakly $\beta(\tau_1, \tau_2)$ -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y such that $F(x) \cap V \neq \emptyset$, there exists a $\tau_1\tau_2\text{-}\beta$ -open set U containing x such that $F(z) \cap \sigma_1\sigma_2\text{-Cl}(V) \neq \emptyset$ for every $z \in U$;
- (3) *upper weakly (resp. lower weakly) $\beta(\tau_1, \tau_2)$ -continuous* if F has this property at each point of X .

3.1 A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is upper weakly $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V containing $F(x)$.

3.2 A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is lower weakly $\beta(\tau_1, \tau_2)$ -continuous at $x \in X$ if and only if $x \in \tau_1\tau_2\text{-}\beta\text{Int}(F^-(\sigma_1\sigma_2\text{Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V such that $F(x) \cap V \neq \emptyset$.

3.3 For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

- (1) F is upper weakly $\beta(\tau_1, \tau_2)$ -continuous;
- (2) $F^+(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V ;
- (3) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(K)))) \subseteq F^-(K)$ for every $\sigma_1\sigma_2$ -closed set K ;

(4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(B)))) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B of Y ;

(5) $F^+(\sigma_1\sigma_2\text{-Int}(B)) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-Cl}(\sigma_1\sigma_2\text{-Int}(B))))$ for every subset B of Y ;

(6) $F^+(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^+(\sigma_1\sigma_2\text{-Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y ;

(7) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^-(V)) \subseteq F^-(\sigma_1\sigma_2\text{-Cl}(V))$ for every $\sigma_1\sigma_2$ -open set V of Y .

3.4 For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following properties are equivalent:

(1) F is lower weakly $\beta(\tau_1, \tau_2)$ -continuous;

(2) $F^-(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^-(\sigma_1\sigma_2\text{Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V of Y ;

(3) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-Int}(K)))) \subseteq F^+(K)$ for every $\sigma_1\sigma_2$ -closed set K ;


(4) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(\sigma_1\sigma_2\text{-Int}(\sigma_1\sigma_2\text{-Cl}(B)))) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(B))$ for every subset B of Y ;

(5) $F^-(\sigma_1\sigma_2\text{-Int}(B)) \subseteq \tau_1\tau_2\text{-}\beta\text{Cl}(F^-(\sigma_1\sigma_2\text{-Cl}(\sigma_1\sigma_2\text{-Int}(B))))$ for every subset B ;

(6) $F^-(V) \subseteq \tau_1\tau_2\text{-}\beta\text{Int}(F^-(\sigma_1\sigma_2\text{-Cl}(V)))$ for every $\sigma_1\sigma_2$ -open set V ;

(7) $\tau_1\tau_2\text{-}\beta\text{Cl}(F^+(V)) \subseteq F^+(\sigma_1\sigma_2\text{-Cl}(V))$ for every $\sigma_1\sigma_2$ -open set V .

3.5 For a multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$, the following implication hold:



The converse of the implications are not true in general.


5.2 Recommendations

To this end, even though we have found several characterizations presented in this thesis, there is another type of continuity that interests us to investigate. The multifunction is defined as:

A multifunction $F : (X, \tau_1, \tau_2) \rightarrow (Y, \sigma_1, \sigma_2)$ is said to be:

- (1) *upper (τ_1, τ_2) -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y containing $F(x)$, there exists a $\tau_1\tau_2$ -open set U containing x such that $F(U) \subseteq V$;
- (2) *lower (τ_1, τ_2) -continuous* at a point $x \in X$ if for each $\sigma_1\sigma_2$ -open set V of Y such that $F(x) \cap V \neq \emptyset$, there exists a $\tau_1\tau_2$ -open set U containing x such that $F(z) \cap V \neq \emptyset$ for every $z \in U$;
- (3) *upper (resp. lower) (τ_1, τ_2) -continuous* if F has this property at each point of X .

Therefore, there are many interesting questions involving several characterizations of this multifunction yet to be answered. Moreover, the relationships between (τ_1, τ_2) -continuity and $\beta(\tau_1, \tau_2)$ -continuity are noticeable to study in the future. Besides, The condition, which make the converse of the relationships true in general, is a further interesting study.







REFERENCES

พูน ปณ จิต ชีเว

REFERENCES

- [1] N. Levine, "Semi-open sets and semi-continuity in topological spaces," *Amer. Math. Monthly*, vol. 70, no. 1, pp. 36–41, 1963.
- [2] A. Mashhour, "On precontinuous and weak precontinuous mappings," in *Proc. Math. Phys. Soc. Egypt.*, vol. 53, pp. 47–53, 1982.
- [3] D. Andrijević, "Semi-preopen sets," *Mat. Vesnik*, vol. 38, no. 93, pp. 24–32, 1986.
- [4] M. A. El-Monsef, S. El-Deeb, and R. Mahmoud, " β -open sets and β -continuous mappings," *Bull. Fac. Sci. Assiut Univ*, vol. 12, no. 1, pp. 77–90, 1983.
- [5] V. Popa and T. Noiri, "On β -continuous functions," *Real Anal. Exchange*, pp. 544–548, 1992.
- [6] J. Borsik and J. Doboš, "On decompositions of quasicontinuity," *Real Anal. Exchange*, vol. 16, no. 1, pp. 292–305, 1990.
- [7] A. Nasef and T. Noiri, "Some weak forms of almost continuity," *Acta Math. Hung.*, vol. 74, no. 3, pp. 211–219, 1997.
- [8] T. Noiri and V. Popa, "On almost β -continuous functions," *Acta Math. Hung.*, vol. 79, no. 4, pp. 329–339, 1998.
- [9] F. Khedr, S. Al-Areefi, and T. Noiri, "Precontinuity and semi-precontinuity in bitopological spaces," *Indian J. Pure Appl. Math.*, vol. 23, pp. 625–625, 1992.
- [10] V. Popa, T. Noiri, *et al.*, "On upper and lower β -continuous multifunctions," *Real Anal. Exchange*, vol. 22, no. 1, pp. 362–376, 1996.
- [11] T. Noiri and V. Popa, "On upper and lower almost β -continuous multifunctions," *Acta Math. Hung.*, vol. 82, no. 1-2, pp. 57–73, 1999.
- [12] V. Popa and T. Noiri, "On upper and lower weakly β -continuous multifunctions," *Anal. Univ. Sci. Budapest*, vol. 43, pp. 25–48, 2000.

- 
- [13] S. W. Davis, *Topology*. McGraw-Hill, 2005.
- [14] J. Kelly, "Bitopological spaces," *Proc. London Math. Soc.*, vol. 3, no. 1, pp. 71–89, 1963.
- [15] J. R. Munkres, *Topology*, ch. 3, p. 164. Prentice Hall, 2000.
- [16] S. Bose, "Semi open sets, semi continuity and semi open mappings in bitopological spaces," *Bull. Cal. Math. Soc.*, vol. 73, pp. 237–246, 1981.
- [17] G. Banerjee, "On pairwise almost strongly θ -continuous mappings," *Bull. Cal. Math. Soc.*, vol. 79, pp. 314–320, 1987.
- [18] M. Jelic, "A decomposition of pairwise continuity," *J. Inst. Math. and Comp. Sci.(Math. Ser.)*, vol. 3, pp. 25–29, 1990.
- [19] S. Maheshwari and R. Prasad, "Semi-open sets and semi-continuous function in bitopological spaces," *Math. Notae*, vol. 26, pp. 29–37, 1977.
- [20] T. Noiri and V. Popa, "On weakly precontinuous functions in bitopological spaces," *Soochow J. Math.*, vol. 33, no. 1, p. 87, 2007.
- [21] C. Boonpok, C. Viriyapong, and M. Thongmoon, "On upper and lower (τ_1, τ_2) -precontinuous multifunctions," *J. Math. Computer Sci*, vol. 18, pp. 282–293, 2018.
- [22] S. Tahiliani, "On weakly β -continuous functions in bitopological spaces," *Filomat*, vol. 22, no. 1, pp. 77–86, 2008.
- [23] C. Berg, "Espaces topologiques fonctions multivoques," 1959.
- [24] C. H. Dowker, "On countably paracompact spaces," *Canadian Journal of Mathematics*, vol. 3, pp. 219–224, 1951.



BIOGRAPHY



NAME	Miss Kaewta Laprom
DATE OF BIRTH	October 26, 1994
PLACE OF BIRTH	Saraburi Province, Thailand
ADDRESS	188, Nongplalai sub-district, Mueang Saraburi district, Saraburi province 18000, Thailand kaewta.lap@msu.ac.th
EDUCATION	
2012	High School in Suankularb Wittayalai Saraburi School, Saraburi, Thailand
2017	Bachelor of Education in Mathematics Education Mahasarakham University, Thailand
2020	Master of Science in Mathematics Education, Mahasarakham University, Thailand